



## Notice of Meeting and Meeting Agenda Core Area Liquid Waste Management Committee

---

Wednesday, March 26, 2025

11:30 AM

6th Floor Boardroom  
625 Fisgard Street  
Victoria, BC V8W 1R7

---

C. Coleman (Chair), D. Kobayashi (Vice Chair), M. Alto, S. Brice, J. Brownoff, J. Caradonna, Z. de Vries, B. Desjardins, S. Goodmanson, C. Harder, K. Murdoch, D. Murdock, C. Plant, D. Thompson, S. Tobias, C. McNeil-Smith (Board Chair, ex-officio)

The Capital Regional District strives to be a place where inclusion is paramount and all people are treated with dignity. We pledge to make our meetings a place where all feel welcome and respected.

### 1. Territorial Acknowledgement

### 2. Approval of Agenda

### 3. Adoption of Minutes

#### 3.1. [25-0370](#) Minutes of the June 26, October 9, December 11, 2024 and January 8, 2025 Core Area Liquid Waste Management Committee Meeting

**Recommendation:** That the minutes of the Core Area Liquid Waste Management Committee meetings of June 26, October 9, and December 11, 2024, and January 8, 2025, be adopted as circulated.

**Attachments:** [Minutes - June 26, 2024](#)  
[Minutes - October 9, 2024](#)  
[Minutes - December 11, 2024](#)  
[Minutes - January 8, 2025](#)

### 4. Chair's Remarks

### 5. Presentations/Delegations

*The public are welcome to attend CRD meetings in-person.*

*Delegations will have the option to participate electronically. Please complete the online application at [www.crd.ca/address](http://www.crd.ca/address) no later than 4:30 pm two days before the meeting and staff will respond with details.*

*Alternatively, you may email your comments on an agenda item to the CRD Board at [crdboard@crd.bc.ca](mailto:crdboard@crd.bc.ca).*

### 6. Committee Business

- 6.1.      [25-0070](#)      2025 Core Area Liquid Waste Management Committee Terms of Reference
- Recommendation:**      There is no recommendation. This report is for information only.
- Attachments:**      [Staff Report: 2025 Core Area Liquid Waste Management ToR](#)  
                                      [Appendix A: 2025 CALWMC Terms of Reference - Approved Jan 8, 2025](#)  
                                      [Appendix B: 2025 CALWMC Terms of Reference - Redlined](#)
- 6.2.      [25-0371](#)      Core Area Wastewater Treatment Plant Odour Mitigation Update
- Recommendation:**      There is no recommendation. This report is for information only.
- Attachments:**      [Staff Report: Core Area WWTP Odour Mitigation Update](#)
- 6.3.      [25-0146](#)      Biosolids Literature Review Outcomes
- Recommendation:**      [At the March 12, 2025 CRD Board meeting, the Board endorsed the following referral motion from the February 19, 2025 Environmental Services Committee meeting: That the Biosolids Literature Review Outcomes report be referred to the Core Area Liquid Waste Management Committee for information.]  
                                      There is no recommendation. This report is for information only.
- Attachments:**      [Staff Report: Biosolids Literature Review Outcomes](#)  
                                      [Appendix A: Biosolids Land Application – Updated Review](#)

## 7. Notice(s) of Motion

## 8. New Business

## 9. Adjournment

The next meeting is May 28, 2025.

To ensure quorum, please advise Jessica Dorman (jdorman@crd.bc.ca) if you or your alternate cannot attend.

## Meeting Minutes

### Core Area Liquid Waste Management Committee

---

Wednesday, June 26, 2024

1:30 PM

6th Floor Boardroom  
625 Fisgard Street  
Victoria, BC V8W 1R7

---

#### PRESENT

Directors: C. Coleman (Chair), D. Kobayashi (Vice Chair), M. Alto, K. Armour (for B. Desjardins), S. Brice, J. Brownoff (EP), J. Caradonna, S. Goodmanson, C. Harder (for L. Szpak), K. Murdoch, D. Murdock (1:34 pm) (EP), C. Plant, D. Thompson (1:31 pm), S. Tobias (EP)

Staff: T. Robbins, Chief Administrative Officer; A. Fraser, General Manager, Integrated Water Services; L. Jones, General Manager, Parks, Recreation and Environmental Services; J. Dales, Senior Manager, Wastewater Infrastructure Operations; J. Marr, Senior Manager, Infrastructure Engineering; J. Kelly, Manager, Integrated Water Services Capital Projects; P. Kickham, Manager, Environmental Regulations; M. Lagoa, Deputy Corporate Officer; J. Dorman, Committee Clerk (Recorder)

EP - Electronic Participation

Regrets: Director(s) Z. de Vries, B. Desjardins, L. Szpak

The meeting was called to order at 1:30 pm.

#### 1. Territorial Acknowledgement

Vice Chair Kobayashi provided a Territorial Acknowledgement.

#### 2. Approval of Agenda

Director Thompson joined the meeting in person at 1:31 pm.

**MOVED** by Director Murdoch, **SECONDED** by Director Kobayashi,  
That the agenda for the June 26, 2024 Core Area Liquid Waste Management  
Committee meeting be approved.

**CARRIED**

#### 3. Adoption of Minutes

##### 3.1. [24-649](#)

Minutes of the February 28, 2024 Core Area Liquid Waste Management  
Committee Meeting

**MOVED** by Director Murdoch, **SECONDED** by Director Plant,  
That the minutes of the Core Area Liquid Waste Management Committee  
meeting of February 28, 2024 be adopted as circulated.

**CARRIED**

#### 4. Chair's Remarks

Chair Coleman and Vice Chair Kobayashi spoke about the map of treatment plants from agenda Item 6.1. and future tours of selected treatment plants.

Director Murdock joined the meeting electronically at 1:34 pm.

#### 5. Presentations/Delegations

There were no presentations or delegations.

#### 6. Committee Business

6.1. [24-625](#) Core Area Liquid Waste Management Committee 2024 Mid-Year Capital Projects and Operations Update

J. Dales and J. Marr presented Item 6.1. for information.

Discussion ensued on the following:

- reduction of odour complaints
- system upgrades and non-compliance events
- plant maintenance and capital projects
- capital budget, cost escalation and fluctuation
- peak capacity and major weather events

6.2. [24-626](#) Core Area Wastewater Treatment Plant Capacity Allocation Review

J. Marr presented Item 6.2. for information

Discussion ensued on sewer upgrades and reallocation of resources.

6.3. [24-570](#) Long-Term Biosolids Management Strategy

P. Kickham presented Item 6.3. for information.

6.4. [24-605](#) Previous Minutes of Other CRD Committees and Commissions for Information

Discussion ensued on the biosolids review and recommendation of the plan.

The following minutes were received for information:

- a) Technical and Community Advisory Committee minutes of February 13, 2024
- b) Technical and Community Advisory Committee minutes of March 14, 2024
- c) Technical and Community Advisory Committee minutes of May 22, 2024

#### 7. Notice(s) of Motion

There were no notice(s) of motion.

**8. New Business**

There was no new business.

**9. Adjournment**

**MOVED** by Director Kobayashi, **SECONDED** by Director Goodmanson,  
That the June 26, 2024 Core Area Liquid Waste Management Committee be  
adjourned at 2:12 pm.  
**CARRIED**

---

CHAIR

---

RECORDER

## Meeting Minutes

### Core Area Liquid Waste Management Committee

---

Wednesday, October 9, 2024

9:00 AM

6th Floor Boardroom  
625 Fisgard Street  
Victoria, BC V8W 1R7

---

#### Special

#### PRESENT

Directors: C. Coleman (Chair), D. Kobayashi (Vice Chair), S. Brice, J. Brownoff (EP), J. Caradonna, Z. de Vries (EP), B. Desjardins, S. Goodmanson (9:03 am) (EP), S. Kim (for M. Alto) (EP), K. Murdoch (EP), D. Murdock (EP), C. Plant, J. Rogers (for S. Tobias) (EP), L. Szpak, D. Thompson

Staff: T. Robbins, Chief Administrative Officer; N. Chan, Chief Financial Officer; A. Fraser, General Manager, Integrated Water Services; J. Dales, Senior Manager, Wastewater Infrastructure Operations; G. Harris, Senior Manager, Environmental Protection; J. Marr, Senior Manager, Infrastructure Engineering; J. Kelly, Manager, Integrated Water Services Capital Projects; P. Kickham, Manager, Environmental Regulations; Y. Li, Senior Financial Advisor; M. Lagoa, Deputy Corporate Officer; J. Dorman, Committee Clerk (Recorder)

EP - Electronic Participation

Regrets: Director(s) M. Alto, S. Tobias

The meeting was called to order at 9:02 am.

#### 1. Territorial Acknowledgement

Vice Chair Kobayashi provided a Territorial Acknowledgement.

#### 2. Approval of Agenda

**MOVED** by Director Desjardins, **SECONDED** by Director Brice,  
That the agenda for the October 9, 2024 Core Area Liquid Waste Management  
Committee be approved.  
**CARRIED**

#### 3. Presentations/Delegations

There were no presentations or delegations.

#### 4. Special Meeting Matters

4.1. [24-941](#)

Core Area Wastewater Service 2025 Operating and Capital Budget

A. Fraser spoke to Item 4.1.

Discussion ensued on the following:

- 5-year projections and forecasting
- chemical cost, procurement, alternatives and competitive rates
- Lafarge biosolids processing capacity
- contributions to reserves and change in assumptions to actuals
- debt retirement strategy analysis
- average annual increase from previous years

**MOVED** by Director Szpak, **SECONDED** by Director Caradonna,  
The Core Area Liquid Waste Management Committee recommends the  
Committee of the Whole recommend to the Capital Regional District Board:

1. Approve the 2025 Core Area Wastewater Service operating and capital budgets as presented;
2. Direct staff to balance the 2024 actual revenue and expenses on the transfer to the operating reserve; and
3. Direct staff to update carry forward balances in the 2025 Capital Budget for changes after year end.

**MOVED** by Director Caradonna, **SECONDED** by Director Brice,  
That the main motion be amended in 1. to add the words "subject to revised,  
more conservative estimates for biosolids-related costs." after the words "as  
presented".

**DEFEATED**

Opposed: Brownoff, Coleman, de Vries, Desjardins, Goodmanson, Murdoch,  
Murdock, Rogers, Szpak, Kim, Kobayashi, Plant

The question was called on the main motion.

The Core Area Liquid Waste Management Committee recommends the  
Committee of the Whole recommend to the Capital Regional District Board:

1. Approve the 2025 Core Area Wastewater Service operating and capital budgets as presented;
2. Direct staff to balance the 2024 actual revenue and expenses on the transfer to the operating reserve; and
3. Direct staff to update carry forward balances in the 2025 Capital Budget for changes after year end.

**CARRIED**

4.2. [24-936](#)

Liquid Waste Management Plan Engagement Plan

G. Harris presented Item 4.2. for information

4.3. [24-957](#) Core Area Wastewater Service Esquimalt Nation Capacity Allocation Request

A. Fraser presented Item 4.3. for information.

Discussion ensued on the correct name of the Nation and CRD's authority versus the process prescribed in the bylaw.

**5. Adjournment**

**MOVED by Director Desjardins, SECONDED by Director Caradonna,  
That the October 9, 2024 Core Area Liquid Waste Management Committee  
meeting be adjourned at 10:40 am.  
CARRIED**

---

CHAIR

---

RECORDER



## Meeting Minutes

### Core Area Liquid Waste Management Committee

---

Wednesday, December 11, 2024

9:00 AM

6th Floor Boardroom  
625 Fisgard Street  
Victoria, BC V8W 1R7

---

#### Special Meeting

##### PRESENT

Directors: C. Coleman (Chair), D. Kobayashi (Vice Chair) (EP), S. Brice, J. Brownoff, J. Caradonna, Z. de Vries (EP), B. Desjardins, S. Goodmanson (EP), C. Harder, S. Kim (for M. Alto), K. Murdoch, D. Murdock (EP), C. Plant, D. Thompson (EP), S. Tobias (EP), C. McNeil-Smith (Ex-Officio)

Staff: T. Robbins, Chief Administrative Officer; A. Fraser, General Manager, Integrated Water Services; G. Harris, Acting General Manager, Parks, Recreation and Environmental Services; L. Jones, General Manager, Parks, Recreation and Environmental Services; K. Morley, General Manager, Corporate Services; J. Dales, Senior Manager, Wastewater Infrastructure Operations; S. May, Senior Manager, Facilities Management and Engineering Services; P. Kickham, Acting Senior Manager, Environmental Protection; M. Lagoa, Deputy Corporate Officer; J. Dorman, Committee Clerk (Recorder)

EP - Electronic Participation

Regrets: Director Alto

The meeting was called to order at 9:02 am.

#### 1. Territorial Acknowledgement

Director Brownoff provided a Territorial Acknowledgement.

#### 2. Approval of Agenda

**MOVED** by Director Brownoff, **SECONDED** by Director Caradonna,  
That the agenda for the December 11, 2024 Core Area Liquid Waste Management  
Committee meeting be approved.  
**CARRIED**

#### 3. Presentations/Delegations

There were no presentations or delegations.

#### 4. Special Meeting Matters

There were no special meeting matters.

## 5. Motion to Close the Meeting

### 5.1. [24-1317](#) Motion to Close the Meeting

**MOVED** by Director Desjardins, **SECONDED** by Director Brice,

**1. That the meeting be closed for Proposed Service in accordance with Section 90(1)(k) of the Community Chater.**

**CARRIED**

**MOVED** by Director Desjardins, **SECONDED** by Director Brice,

**2. That such disclosures could reasonably be expected to harm the interests of the Regional District.**

**CARRIED**

The Core Area Liquid Waste Management Committee moved into the closed session at 9:05 am.

The Core Area Liquid Waste Management Committee rose from the closed session at 10:52 am without report.

## 6. Adjournment

**MOVED** by Director Browhoff, **SECONDED** by Director Murdoch,

**That the December 11, 2024 Core Area Liquid Waste Management Committee be adjourned at 10:52 am.**

**CARRIED**

---

CHAIR

---

RECORDER

## Meeting Minutes

### Core Area Liquid Waste Management Committee

---

Wednesday, January 8, 2025

9:00 AM

6th Floor Boardroom  
625 Fisgard Street  
Victoria, BC V8W 1R7

---

#### Special Meeting

#### PRESENT

Directors: C. Coleman (Chair), D. Kobayashi (Vice Chair), S. Brice, J. Brownoff, J. Caradonna, Z. de Vries, B. Desjardins, S. Goodmanson, C. Harder (9:04 am), S. Kim (for M. Alto) (EP), K. Murdoch, D. Murdock, D. Thompson, S. Tobias, C. McNeil-Smith (Ex-Officio)

Staff: T. Robbins, Chief Administrative Officer; N. Chan, Chief Financial Officer; A. Fraser, General Manager, Infrastructure & Water Services; L. Jones, General Manager, Parks, Recreation and Environmental Services; K. Morley, General Manager, Corporate Services; G. Harris, Senior Manager, Parks, Recreation and Environmental Services; S. May, Senior Manager, Facilities Management and Engineering Services; P. Kickham, Manager, Environmental Regulations; M. Lagoa, Deputy Corporate Officer; J. Dorman, Committee Clerk (Recorder)

EP - Electronic Participation

Regrets: M. Alto, C. Plant

The meeting was called to order at 9:00 am.

#### 1. Territorial Acknowledgement

Vice Chair Kobayashi provided a Territorial Acknowledgement.

#### 2. Approval of Agenda

**MOVED by Director Murdoch, SECONDED by Director Kobayashi,  
That the January 8, 2025 Core Area Liquid Waste Management Committee  
agenda be approved as amended with the addition of the following item:**

**- 3. Chair's Remarks**

**CARRIED**

#### 3. Chair's Remarks

Chair Coleman spoke about the opportunity for Directors to participate in future orientation tours of the treatment plants.

#### 4. Presentations/Delegations

There were no presentations or delegations.

## 5. Special Meeting Matters

There were no Special Meeting Matters.

## 6. Motion to Close the Meeting

### 6.1. [24-1359](#) Motion to Close the Meeting

**MOVED** by Director Kobayashi, **SECONDED** by Director Murdoch,  
That the meeting be closed for Proposed Service in accordance with Section (90)  
(1)(k) of the Community Charter.

**CARRIED**

**MOVED** by Director Kobayashi, **SECONDED** by Director Murdoch,  
That such disclosures could reasonably be expected to harm the interests of the  
Regional District.

**CARRIED**

Director Harder joined the meeting at 9:04 am.

The Core Area Liquid Waste Management Committee went into closed session  
at 9:04 am.

The Core Area Liquid Waste Management Committee rose from closed  
session at 11:54 am without report.

## 7. Adjournment

**MOVED** by Director Murdoch, **SECONDED** by Director Murdoch,  
That the January 8, 2025 Core Area Liquid Waste Management Committee  
meeting be adjourned at 11:54 am.

**CARRIED**

---

CHAIR

---

RECORDER

**REPORT TO CORE AREA LIQUID WASTE MANAGEMENT COMMITTEE  
MEETING OF WEDNESDAY, MARCH 26, 2025**

---

**SUBJECT**     **2025 Core Area Liquid Waste Management Committee Terms of Reference**

**ISSUE SUMMARY**

To provide the 2025 Core Area Liquid Waste Management Committee Terms of Reference for information.

**BACKGROUND**

Under the *Local Government Act* and the CRD Board Procedures Bylaw, the CRD Board Chair has the authority to establish standing committees and appoint members to provide advice and recommendations to the Board.

On January 8, 2025, the CRD Board approved the 2025 Terms of Reference for standing committees. Terms of Reference (TOR) serve to clarify the mandate, responsibilities and procedures of standing committees and provide a point of reference and guidance for the committees and members.

The Core Area Liquid Waste Management Committee TOR was updated to reflect recent changes to the General Manager titles as a result of CRD Evolves.

The approved 2025 Core Area Liquid Waste Management Committee TOR is attached as Appendix A, and a redlined copy is attached as Appendix B.

The TOR are being provided for information to the Committee. Any proposed revisions to the TOR will require ratification by the Board.

**CONCLUSION**

Terms of Reference serve to clarify the mandate, responsibilities and procedures of committees and provide a point of reference and guidance for the committees and their members. Any future revisions to the TOR will require ratification by the Board.

**RECOMMENDATION**

There is no recommendation. This report is for information only.

Submitted by:	Marlene Lagoa, MPA, Manager, Legislative Services & Deputy Corporate Officer
Concurrence:	Alicia Fraser, P. Eng., General Manager, Infrastructure and Water Services
Concurrence:	Luisa Jones, MBA, General Manager, Parks, Recreation & Environmental Services
Concurrence:	Kristen Morley, J.D., General Manager, Corporate Services & Corporate Officer
Concurrence:	Ted Robbins, B. Sc., C. Tech., Chief Administrative Officer

**ATTACHMENT(S)**

Appendix A: 2025 Core Area Liquid Waste Management Committee TOR – Approved  
Appendix B: 2025 Core Area Liquid Waste Management Committee TOR – Redlined



## **CORE AREA LIQUID WASTE MANAGEMENT COMMITTEE**

### **PREAMBLE**

The Capital Regional District (CRD) Core Area Liquid Waste Management Committee (CALWMC) is a standing committee established by the CRD Board and will oversee and make recommendations to the Board regarding the Core Area Liquid Waste Management Plan (CALWMP). Recommendations related to long-term biosolids management planning in the CALWMP shall be referred to the Environmental Services Committee.

The Committee's official name is to be:

Core Area Liquid Waste Management Committee

### **1.0 PURPOSE**

- a) The mandate of the Committee is to oversee and make recommendations to the Board regarding the:
  - i. administration and regulatory reporting for the Core Area Liquid Waste Management Plan
  - ii. Core area trunk sewers and sewage disposal systems
- b) The Committee will act as the steering committee of the Technical and Community Advisory Committee, as outlined in Appendix A.

### **2.0 ESTABLISHMENT AND AUTHORITY**

- a) The Committee will make recommendations to the Board for consideration.
- b) The Board Chair will appoint the Committee Chair, Vice Chair and Committee members annually.

### **3.0 COMPOSITION**

- a) The membership is comprised of all directors on the CRD Board from the following municipalities that are participants in the Core Area Liquid Waste Management Plan:
  - Colwood
  - Esquimalt
  - Langford
  - Oak Bay
  - Saanich
  - Victoria
  - View Royal
  - An elected representative and alternate from each of the Songhees Nation and Esquimalt Nation Councils (Board Procedures Bylaw No. 3828)

- b) All Board members are permitted to participate in standing committee meetings, but not vote, in accordance with the CRD Board Procedures Bylaw; and
- c) First Nation members are permitted to participate in standing committee meetings at their pleasure, in accordance with the CRD Procedures Bylaw, where the Nation has an interest in matters being considered by the committee.

#### **4.0 PROCEDURES**

- a) The Committee shall meet quarterly and have special meetings as required at the call of the Committee Chair;
- b) The agenda will be finalized in consultation between staff and the Committee Chair and any Committee member may make a request to the Chair to place a matter on the agenda through the Notice of Motion process;
- c) With the approval of the Committee Chair and Board Chair, Committee matters of an urgent or time sensitive nature may be forwarded directly to the Board for consideration; and
- d) A quorum is a majority of the Committee membership and is required to conduct Committee business.

#### **5.0 RESOURCES AND SUPPORT**

- a) The General Managers of Infrastructure and Water Services, and Parks, Recreation and Environmental Services will act as a liaison to the Committee with support from other departments, as required; and
- b) Minutes and agendas are prepared and distributed by the Corporate Services Department.

*Approved by CRD Board January 8, 2025*



**APPENDIX A**

**STEERING THE TECHNICAL AND COMMUNITY ADVISORY COMMITTEE**

In accordance with the Terms of Reference of the Technical and Community Advisory Committee (TCAC) approved by the Capital Regional District Board (CRD), October 11, 2023, the Core Area Liquid Waste Management Committee (CALWMC) will steer the TCAC as follows:

- Make requests to TCAC for appropriate technical and community consultation advice and input in order to facilitate informed decision-making in a variety of CALWMP matters
- Dissolve the TCAC at a time determined by the CALWMC



## **CORE AREA LIQUID WASTE MANAGEMENT COMMITTEE**

### **PREAMBLE**

The Capital Regional District (CRD) Core Area Liquid Waste Management Committee (CALWMC) is a standing committee established by the CRD Board and will oversee and make recommendations to the Board regarding the Core Area Liquid Waste Management Plan (CALWMP). Recommendations related to long-term biosolids management planning in the CALWMP shall be referred to the Environmental Services Committee.

The Committee's official name is to be:

Core Area Liquid Waste Management Committee

### **1.0 PURPOSE**

- a) The mandate of the Committee is to oversee and make recommendations to the Board regarding the:
  - i. administration and regulatory reporting for the Core Area Liquid Waste Management Plan
  - ii. Core area trunk sewers and sewage disposal systems
- b) The Committee will act as the steering committee of the Technical and Community Advisory Committee, as outlined in Appendix A.

### **2.0 ESTABLISHMENT AND AUTHORITY**

- a) The Committee will make recommendations to the Board for consideration.
- b) The Board Chair will appoint the Committee Chair, Vice Chair and Committee members annually.

### **3.0 COMPOSITION**

- a) The membership is comprised of all directors on the CRD Board from the following municipalities that are participants in the Core Area Liquid Waste Management Plan:
  - Colwood
  - Esquimalt
  - Langford
  - Oak Bay
  - Saanich
  - Victoria
  - View Royal
  - An elected representative and alternate from each of the Songhees Nation and Esquimalt Nation Councils (Board Procedures Bylaw No. 3828)

- b) All Board members are permitted to participate in standing committee meetings, but not vote, in accordance with the CRD Board Procedures Bylaw; and
- c) First Nation members are permitted to participate in standing committee meetings at their pleasure, in accordance with the CRD Procedures Bylaw, where the Nation has an interest in matters being considered by the committee.

#### 4.0 PROCEDURES

- a) The Committee shall meet quarterly and have special meetings as required at the call of the Committee Chair;
- b) The agenda will be finalized in consultation between staff and the Committee Chair and any Committee member may make a request to the Chair to place a matter on the agenda through the Notice of Motion process;
- c) With the approval of the Committee Chair and Board Chair, Committee matters of an urgent or time sensitive nature may be forwarded directly to the Board for consideration; and
- d) A quorum is a majority of the Committee membership and is required to conduct Committee business.

#### 5.0 RESOURCES AND SUPPORT

- a) The General Managers ~~of, Infrastructure and Integrated~~ Water Services, and ~~General Manager, Parks, Recreation and~~ Environmental Services will act as a liaison to the Committee with support from other departments, as required; and
- b) Minutes and agendas are prepared and distributed by the Corporate Services Department.

Approved by CRD Board \_\_\_\_\_

**APPENDIX A**

**STEERING THE TECHNICAL AND COMMUNITY ADVISORY COMMITTEE**

In accordance with the Terms of Reference of the Technical and Community Advisory Committee (TCAC) approved by the Capital Regional District Board (CRD), October 11, 2023, the Core Area Liquid Waste Management Committee (CALWMC) will steer the TCAC as follows:

- Make requests to TCAC for appropriate technical and community consultation advice and input in order to facilitate informed decision-making in a variety of CALWMP matters
- Dissolve the TCAC at a time determined by the CALWMC



Making a difference...together

## **REPORT TO CORE AREA LIQUID WASTE MANAGEMENT COMMITTEE MEETING OF WEDNESDAY, MARCH 26, 2025**

---

**SUBJECT**     **Core Area Wastewater Treatment Plant Odour Mitigation Update**

### **ISSUE SUMMARY**

To provide an update on the Core Area Odour Mitigation Strategy, a summary of actions taken and planned projects moving forward.

### **BACKGROUND**

The McLoughlin Point Wastewater Treatment Plant (MPWWTP) was designed in accordance with the Project Agreement, which included treatment of all odour-laden air before discharge to the environment. All exhausted air at and beyond the plant boundary property line and the shoreline should have a maximum odour concentration of five odour units per cubic meter (OU/m<sup>3</sup>). MPWWTP has a sophisticated air collection and handling system, with carbon treatment to manage odour-laden air from the process systems. Despite the state-of-the-art odour control systems, MPWWTP has generated complaints from neighbouring communities since commissioning. As a result, the Capital Regional District (CRD) committed to continued monitoring of the odour control system performance and discharges, and to work directly with equipment suppliers to seek opportunities to further increase potential odour removal.

An internal audit of the odour collection and treatment systems, as well as an analysis of the previous odour complaints, completed at the end of 2022, outlined the odour mitigation steps for future years. These tasks included an odour assessment using dispersion modelling, odour collection upgrades, revisions to plant maintenance procedures, and a recommendation for further review of findings to validate the approach.

Results from the odour modelling work completed by WSP Canada Inc. confirmed the maximum odour concentration at the plant property boundary remains below five OU/m<sup>3</sup> during normal plant operations. During plant maintenance work, when tank covers are removed, odour concentrations at the plant property boundary and within the surrounding community were observed to exceed the maximum five OU/m<sup>3</sup>. Staff continue to log and investigate all odour complaints from the community. This information is used to further inform the modelling and understanding of the impact of maintenance activities on odour dispersion.

### Odour Complaints and Public Engagement

In 2024, there were 73 odour complaints received, and each odour complaint was investigated to establish the nature of the odour.

Table 1: Summary of Wastewater Odour Complaints Received

<b>Year</b>	<b>Total Odour Incidents</b>	<b>Number of unique complaints</b>
<b>2018</b>	11	11
<b>2019</b>	19	19
<b>2020</b>	29	27
<b>2021</b>	222	69
<b>2022</b>	101	29
<b>2023</b>	149	46
<b>2024</b>	73	33

Odour complaints were received from CRD residents across the region for concerns related to infrastructure in the Core Area, Saanich Peninsula, Southern Gulf Islands, and at least three directly attributed to non-CRD infrastructure. Historically, several odour complaints were attributed to 16 identified cross connections within the City of Victoria collection system. In 2024, the City of Victoria undertook repairs in Barnard Park eliminating cross connections and mitigating a significant odour source.

Of the 73 odour complaints in 2024, staff estimate that 62 were attributed to the Core Area system. Of those 62 complaints, 49 were directly attributed to MPWWTP and 13 attributed to the conveyance system.

Staff continue to engage the Esquimalt Liaison Committee (ELC) on the odour mitigation strategy and ongoing odour concerns. In addition, the ELC is proactively informed of potential odour generating maintenance work. In 2025 semi-annual meetings are scheduled with this committee (February 27 and September 25), with two operational updates provided May 22 and November 27.

### Administrative Improvements

As maintenance activities have been identified as the primary source of odour exceedances outside of the plant property boundary, staff continue to refine and optimize the scheduling and execution of maintenance activities to minimize potential odours.

- Standard operating procedures now require a review of these factors to align potential odour-causing maintenance activities with more favorable atmospheric conditions.
- Improved procedures focused on reducing the required time covers are removed. When covers must be removed, operators track odour conditions and increase the application of an odour neutralizing compound (Ecosorb), now aided with an industrial dispersion fan.

The drop in complaints received in 2024 has been aided by administrative improvements and is projected to improve as numerous equipment upgrades are scheduled to complete later in 2025.

### 2025 Backwash Tank Maintenance

Through the treatment process at MPWWTP, the dirty backwash tank slowly accumulates a granular material used in the biological aerated filters. The design of the plant requires maintenance on a 4–5-year cycle to ensure efficient treatment operation and reduce the risk of odour generation. To access this infrastructure, CRD staff will need to bypass secondary treatment at the facility, resulting in effluent that has undergone primary treatment exiting the outfall into the Salish Sea for 21 days. Permission to proceed with the project required both federal and provincial approval, including notification to First Nations and the public. This will be the largest maintenance project since commissioning in January 2021 and is set to begin on March 26, 2025. For eight days starting March 29, 2025 staff anticipate the potential for increased localized odour generation while staff and contractors work continuously day and night. During this scheduled maintenance, CRD staff will utilize every administrative improvement developed over the last three years to mitigate the odour impact.

### 2025 System Upgrades

Infrastructure work scheduled for 2025, to improve the system performance and reduce odour emissions, includes:

- Secondary Odour Control System Upgrades: Reduced moisture through the pre-filter improves odour elimination and extends the lifespan of the product and reduces maintenance activities.
- Densadeg No. 1 Scum Removal System: Improved scum removal eliminates an odour source and reduces the required frequency of Plant Maintenance.
- MBBR Odour Extraction Ducting Improvements: The addition of mist eliminators and increasing the duct diameter will improve odour extraction from the MBBR tank.
- Dirty Backwash Tank Odour Treatment System Upgrade: Upgrading the current passive system to a larger fan unit.

### Future 2026 Improvements

- Collection system full-service odour control system.
- Upgrades to the primary odour control system mist eliminator.
- Additional odour extraction points for plate settlers and Densadegs.

### **CONCLUSION**

The Capital Regional District recognizes that while operating under normal conditions the McLoughlin Point Wastewater Treatment Plant (MPWWTP) odour control system is fully functional and operating as per design specifications; however, additional system improvements are being made, according to the MPWWTP Odour Control System Review and Assessment. In 2025 and 2026, this odour mitigation project work will continue to improve regular plant operation and reduce the community odour impact from plant maintenance.

**RECOMMENDATION**

There is no recommendation, this report is for information only.

Submitted by:	Jason Dales, B.SC., WD IV, Senior Manager, Wastewater Infrastructure Operations
Concurrence:	Alicia Fraser, P. Eng., General Manager, Infrastructure and Water Services
Concurrence:	Ted Robbins, B. Sc., C. Tech., Chief Administrative Officer



**REPORT TO ENVIRONMENTAL SERVICES COMMITTEE  
MEETING OF WEDNESDAY, FEBRUARY 19, 2025**

---

**SUBJECT     Biosolids Literature Review Outcomes**

**ISSUE SUMMARY**

To present the Environmental Services Committee with the results of the independent academic literature review on the risks and benefits of biosolids land application.

**BACKGROUND**

At the August 9, 2023 Capital Regional District (CRD) Board meeting, staff were directed to *report back with a proposal that CRD Environment Service fund University of Victoria or other suitable independent academic institution to prepare a review: a) of available literature, to determine whether there are validated examples and/or peer reviewed papers assessing the risks and benefits of the application of biosolids on environmental and human health, and b) based on this and on The Precautionary Principle, whether CRD may have a legal liability for such application.*

At the October 18, 2023, Environmental Services Committee meeting, staff presented a proposal for an academic institution to conduct a literature review. At that time, the provincial government's Technical Working Group (TWG) was expected to issue a report on its review of the Organic Matter Recycling Regulation in late 2023. Given the upcoming report, the committee passed the following motion: *That the committee postpone discussion on this item until the January 2024 committee meeting.*

After delays to the release of the TWG report, the CRD Board directed staff to reinstitute the process of authorizing the literature review by the following motion at the March 13, 2024 Board meeting: *Given delays to provincial reporting on Organic Matter Recycling Regulation, and the Board's previous direction to initiate an academic analysis, that the Board direct staff to move forward with a third-party academic review of the scientific literature on the uses and impacts of biosolids.*

At the June 19, 2024 Environmental Services Committee meeting, staff presented qualification criteria for a suitable academic researcher. The Committee directed staff to 1. *Secure a tenured professor that fulfills the qualification criteria outlined in this report, to undertake the independent literature review, as per the terms of reference previously approved for this work, with a budget not to exceed \$40,000; and 2. That staff be directed to procure a legal review in alignment with the selection criteria and scope of work presented in this report, with a budget not to exceed \$25,000.*

Dr. Chris Kennedy, a Professor of Aquatic Toxicology at Simon Fraser University was selected to author the literature review. The report (Appendix A), focused on relevant literature published in 2023 and 2024; the work was completed in late 2024.

The law firm Borden Ladner and Gervais is completing the review of legal liability associated with land application of biosolids. The review is in the late stage of drafting, and staff anticipate providing the report to the ESC in March.

The objectives of the literature review were to provide up to date information on the following:

1. The human health and environmental risks of both legacy contaminants and Contaminants of Emerging Concern (COECs), with consideration of environmental conditions typical of the BC south coast.
2. Contaminant concentrations in biosolids relative to levels of exposure in general society.
3. The limitations of extrapolating lab-based toxicity testing to observations in the environment.
4. A summary of the areas of uncertainty in biosolids land application risk, including a summary of relevant techniques for evaluating and addressing uncertainty.
5. A summary of biosolids land application techniques that can reduce risk and/or address uncertainty.
6. A summary of the risks and concerns that have resulted in land application bans elsewhere.
7. An assessment of the overall risks of biosolids land application considering the intent of the Precautionary Principle (Rio Declaration, 1992 and subsequent derivations).

The literature review considered a broad suite of COECs including per and polyfluoroalkyl substances (PFAS), pharmaceuticals and personal care products and microplastics.

## **IMPLICATIONS**

### *Environmental Implications*

The author concluded that the low concentration of COECs in CRD biosolids coupled with existing toxicity data suggests that the COECs in CRD biosolids represent a negligible to low risk to human health and the environment. However, the report also highlighted significant sources of uncertainty in this conclusion, which should be reviewed regularly as the science evolves.

An adaptive management framework is recommended to mitigate uncertainty by regularly reviewing available updates to the scientific literature. Under such a framework, decision makers keep pace with evolving science and regularly weigh the risks and benefits in biosolids management. In the context of biosolids land application, this means enhanced monitoring of COECs in biosolids, careful site selection, and the adoption of advanced treatment technologies as they become available.

### *Alignment with Existing Plans & Strategies*

The conclusions of the literature review align with the CRD's proposed Long-term Biosolids Management Strategy (long-term strategy), as Tier 1 of the long-term strategy includes investigation and development of emerging technology (advanced thermal demonstration facility) for biosolids management. The long-term strategy also exercises a precautionary approach by excluding agricultural land application options. Staff have also begun testing biosolids biannually for a greater range of contaminants, including PFAS and pharmaceuticals, which can inform an adaptive management framework where staff will continue to monitor the evolving science to inform decision-making.

## **CONCLUSION**

A literature review commissioned by the CRD examined the human health and environmental risks and benefits of biosolids land application. Based on recent research, the review concluded

that contaminants of emerging concern (COECs) present in CRD biosolids pose a negligible to low risk to human health and the environment. However, it also identified significant sources of uncertainty in this risk assessment. To address this uncertainty, the review recommended an adaptive management framework that includes exercising caution and exploring emerging technologies. The CRD's proposed long-term strategy aligns with these conclusions by prioritizing the development of an advanced thermal demonstration facility and excluding agricultural land application from consideration. Staff will continue to monitor evolving science and have initiated enhanced monitoring of COECs in biosolids to inform future decision-making.

### **RECOMMENDATION**

There is no recommendation. This report is for information only.

Submitted by:	Peter Kickham, M.E.T., R.P.Bio., Acting Senior Manager, Environmental Innovation & Strategy
Concurrence:	Luisa Jones, MBA, General Manager, Parks, Recreation & Environmental Services
Concurrence:	Ted Robbins, B. Sc., C. Tech., Chief Administrative Officer

### **ATTACHMENT**

Appendix A: Biosolids Land Application – An Updated Review of Human Health and Environmental Risks

# FINAL REPORT

## Biosolids Land Application – An Updated Review of Human Health and Environmental Risks

Version 2.0

December 12, 2024

Prepared for:

Capital Regional District

Prepared by:

Biowest Environmental Research Consultants

Dr. Christopher J. Kennedy

# Table of Contents

1	Introduction	1
2	Contaminants Present in Biosolids and Associated Risks	4
2.1	PFAS	5
2.1.1	Screening Level Human Health and Ecological Risk Assessment for PFAS in Biosolids	6
2.1.2	Estimated Daily Intakes of PFOS and PFOA	8
2.2	Microplastics	9
2.3	Pharmaceuticals and personal care products	10
2.4	Prioritization of Unregulated Organic Chemicals in Biosolids (Warke and McAvoy, 2024)	12
3	Fate and Transport of Contaminants in Biosolids and Considerations of Conditions Typical of the BC South Coast	13
4	Techniques to Reduce Risks in Biosolids Land Application	15
4.1	Recent Research on Treatment Technologies	15
4.2	Application Techniques and Site-Specific Application Considerations	16
5	Limitations of Extrapolating Lab-Based Testing to the Environment	17
6	Uncertainties in the Risks Associated with the Land Application of Biosolids	19
7	Biosolid Land Application Bans	21
7.1	United States of America	21
7.2	Europe	21
8	Discussion and Conclusion	22
9	Limitations	23
10	References	24

# 1 Introduction

Municipal wastewater treatment yields two primary byproducts: treated wastewater and solid residue, or biosolids. Soil application of biosolids as an organic fertilizer is an economic way to beneficially use the carbon and nutrient contents in biosolids to promote soil fertility. A substantial body of research has explored the risks and benefits associated with the land application of biosolids, including numerous studies focused on contaminants, including heavy metals, pathogens, and contaminants of emerging concern (COECs) (Hydromantis Inc., 2010; McCarthy et al., 2015; LRCS Land Resource Consulting Services, 2016; Pozzebon and Seifert, 2023; BC ENV 2024).

There are many benefits associated with the land application of biosolids. Benefits include the addition of organic matter to soil and the associated increase in soil water retention and reduction in soil erosion, as well as the addition of essential nutrients for plant growth. Biosolids can provide a natural alternative to synthetic fertilizers and can store carbon in soil and reduce greenhouse gas emissions (LRCS Land Resource Consulting Services, 2016; BC ENV, 2024). These benefits have been well documented by others and are not the focus of this review. The objective of this report is to build on previous literature reviews assessing potential human health and environmental risks through a comprehensive scan of the recent primary scientific literature and other relevant studies.

As background, the British Columbia Organic Matter Recycling Regulation (OMRR) (available at <https://www.bclaws.gov.bc.ca/OMRR>) provides a regulatory framework for the production, quality, and application of organic matter products, including biosolids. Its purpose is to ensure that biosolids and other organic materials are used in a way that protects human health and the environment while promoting sustainable resource use. The OMRR was developed based on a review of the available scientific literature in the late 1990s and implemented in 2002. The OMRR has undergone several regulatory reviews, with additional amendments proposed which are noted to include provisions to enable the BC Ministry of Environment and Climate Change Strategy to require sampling and analysis for COECs (BC ENV, 2024). Currently, while the OMRR includes requirements for vector (i.e., rodents, birds and insects) attraction reduction processes, and limits for pathogens and heavy metals, the regulation does not include limits for several key COECs including per- and polyfluoroalkyl substances (PFAS), pharmaceuticals and personal care products (PPCPs), and microplastics. While it is acknowledged that the derivation of limits for some of the COECs is complicated by the limited available toxicological data, the importance of recommending limits based on the best available science is essential for the protection of human health and the environment.

In response to concerns over the land application of biosolids, the BC government has convened two technical working groups (TWGs), one in 2015 and a second in 2022, to review scientific information on biosolids production and waste management practices. A summary of the findings of the reports produced by the two TWGs follows.

The findings of the 2015 Technical Working Group were presented in the report titled *A literature review of risks relevant to the use of biosolids and compost from biosolids with relevance to the Nicola Valley, BC* (LRCS Land Resource Consulting Services, 2016). The review focused on risks not addressed by the existing regulatory framework for biosolids management in Canada, including the issue of COECs. Primary environmental and human health risks identified included the potential contamination of ground and surface water by COECs, seasonal risks due to groundwater recharge and runoff following biosolids application, and the potential for livestock and wildlife to be directly exposed to contaminants in biosolids via ingestion of biosolids during foraging, as well as the ingestion of plants that have accumulated the contaminants. Several knowledge gaps were identified, including:

- Insufficient empirical data to conduct detailed human health and ecological risk

assessments.

- Lack of data on the ever-expanding list of emerging contaminants of concern present in biosolids.
- Lack of research on the potential synergistic effects of contaminants present in biosolids.
- Limited field studies on biosolid impacts on ground and surface water and subsequent effects in aquatic receiving environments.
- Limited studies assessing the effects of the land application of biosolids on wildlife at all trophic levels.
- Gaps in the understanding of the ability of treatment processes to reduce toxic loading to environmental media and biota.

The report presented several recommendations, including implementing routine public reporting on biosolid composition, focusing on contaminants, including COECs, conducting quantitative risk assessments to address area-specific exposures and associated risks, revising regulations to increase setback distances from watercourses for Class B biosolids and enhanced monitoring programs, to further evaluate the effectiveness of composting and thermal treatment to reduce COECs and to develop public education and source reduction programs to reduce the introduction of contaminants into wastewater systems.

The 2022 TWG provided recommendations in a report titled Organic Matter Recycling Regulation, Technical Working Group Report 2024. The scope of the 2022 TWG was limited and focused on identifying new scientific information on biosolids and compost constituents and management, and on new information on COECs in biosolids and compost since the 2015 TWG. BC ENV (2024) provides the following key messages derived from the TWG findings:

- Each compost and biosolids product is unique to their origin. For example, wastewater treatment plants vary in their input sources (e.g., industrial vs. residential) and may use different treatment processes. Further, the land to which the biosolids (and compost) are applied have unique characteristics. As such, an OMRR application plan specific to the source and the application site should be prepared.
- The importance of source control as the quality of the biosolids (and compost) is directly related to their inputs. The 2022 TWG therefore advocated for regulations that focus on preventing contamination at the source.
- Our understanding is constantly evolving as science advances, and it is challenging to keep pace with evolving science. The 2022 TWG therefore recommended that the ministry devote resources to monitoring the scientific literature and be transparent regarding the research that is used to inform policy.
- To identify and manage COECs it was recommended that the BC ENV develop a comprehensive and transparent strategy that should include not only the presence of COECs, but associated risk. The TWG recommended that the ENV focus on the results of field-based, vs. laboratory-based, studies.
- Provision of context for clear, factual and easily understood information, including providing plain language information on the BC ENV website for why we compost and use biosolids.

Given the large body of scientific literature on the benefits and risks of biosolids that exists, as well as the work conducted by the TWGs convened by the BC ENV, this review is focused on the scientific literature and other relevant reports that have been produced over the last two years (2023 and 2024), with the objective to provide up to date information on the following:

- The human health and environmental risks of both legacy contaminants and COECs, with consideration of environmental conditions typical of the BC south coast.
- Contaminant concentrations in biosolids relative to levels of exposure in general society.
- The limitations of extrapolating lab-based toxicity testing to observations in the environment.
- A summary of the areas of uncertainty in biosolids land application risk, including a summary of relevant techniques for evaluating and addressing uncertainty.
- A summary of biosolids land application techniques that can reduce risk and/or address uncertainty.
- A summary of the risks and concerns that have resulted in land application bans elsewhere.
- An assessment of the overall risks of biosolids land application considering the intent of the Precautionary Principle (Rio Declaration, 1992 and subsequent derivations).



## 2 Contaminants Present in Biosolids and Associated Risks

Previous studies have identified the presence of numerous contaminants in biosolids. These include legacy contaminants such as heavy metals (Sloan et al., 1997; Evanylo et al., 2006; LRCS Land Resource Consulting Services, 2016; Marchuk et al., 2023) and persistent organic pollutants like polychlorinated biphenyls (PCBs), dioxins and furans (PCDD/Fs) and polycyclic aromatic hydrocarbons (PAHs) (Furr et al., 1976; Bergh and Peoples, 1977). In addition, several COECs have been identified in biosolids, including PFAS (LRCS Land Resource Consulting Services, 2016; Wang et al., 2020; Moodie et al., 2021), microplastics (Crossman et al., 2020; Mohajerani and Karabatak, 2020), PPCPs (LRCS Land Resource Consulting Services, 2016; Mohajerani and Karabatak, 2020; Kinney and Heuvel, 2020) and industrial contaminants such as plasticizers, surfactants and brominated flame retardants. The USEPA (<https://www.epa.gov/biosolids/basic-information-about-biosolids>) reports that more than 700 contaminants have been identified in biosolids (in at least one instance) since 1993, with the contaminants present in biosolids varying between wastewater treatment plants (WWTPs) depending on inputs to the facilities.

Regulatory agencies, including the BC ENV, have established limits for metals in biosolids, as well as for pathogens. These contaminants are routinely monitored in biosolids prior to land application. Further, as the toxicity of the legacy organic pollutants (e.g., PCBs, PCDD/F, PAHs) is well understood, many of them have existing standards or criteria in Canada and internationally. A sampling program conducted by the BC ENV (2019) indicated that the concentrations of metals, pathogens and legacy organic pollutants were less than the applicable standards/guidelines for biosolids in samples collected from two WWTPs in the province. Further, previous assessments (Smith, 2009; Eriksen et al., 2009; Jensen et al., 2012; WEAO 2001 and 2010; Higgin et al., 2010) have evaluated risks associated with these contaminants in biosolids. These previous studies have indicated a negligible to low potential for risk to human health and the environment. As the use of some of these chemicals have been phased out overtime, concentrations of these contaminants in wastewater (and biosolids) have also decreased overtime. Despite having a good understanding of the potential risks associated with legacy contaminants in biosolids on an individual level, many of the uncertainties discussed throughout this report, including the lack of understanding of the potential for additive and synergistic effects of contaminants in biosolids, also apply to legacy contaminants.

Contaminants of emerging concern are those that have been identified in recent years, primarily due to advancements in laboratory methodologies and the associated reduction in laboratory detection limits. As they are relatively new contaminants, with limited toxicological data, regulatory agencies typically have not derived environmental quality standards for COECs. Further, due to the same limitation as well as a lack of understanding of the fate of these contaminants following land application of biosolids, a limited number of risk assessments have been conducted to assess the potential for COECs in biosolids to adversely impact human health and the environment. Those available (Eriksen, 2009; Smith, 2009; Jensen et al., 2012; Higgins et al., 2010; Kennedy/Jenks, 2017) have concluded that the contaminants generally represent a low risk to human and environmental health based on the low concentrations of most of the COECs identified in biosolids. Further studies (Higgins et al., 2010; Clarke and Smith, 2011; TCEQ , 2021; Warke and McAvoy, 2024) have attempted to prioritize or rank COECs in biosolids, but have encountered similar challenges due to the paucity of toxicity data for many of the COECs.

A review of the recent (2023-2024) literature pertaining to COECs in biosolids indicates that the scientific understanding of select COECs in biosolids is evolving at a rapid rate, with many recent studies focused on PFAS, microplastics and PPCPs. Research focused on the prevalence of these and other COECs in biosolids, their fate following land application, and their toxicities is being undertaken across the globe, including in Canada. This research has highlighted the existing data gaps and uncertainties in our understanding of COECs in biosolids owing to the complex mixture of contaminants present in biosolids,

the variability in the contaminant profile observed between sources depending on inputs, and the potential for these contaminants to act additively or synergistically.

With the objective of this report to update previous studies regarding human health and ecological risks from the land application of biosolids, a summary of the recent literature and other information (e.g., regulatory decisions) pertaining to PFAS, microplastics and PPCPs follows. Recent data on the concentrations of PFAS in Canadian biosolids was identified, including from the Capital Regional District (CRD), and has been used to conduct a screening level human health and ecological risk assessment of select PFAS with available toxicological data (Section 2.1.1), as well as a comparison of exposures of PFOA and PFOS in CRD biosolids relative to typical daily exposures intakes of PFOA and PFOS (Section 2.1.2). Additionally, a summary of a 2024 study conducted to rank unregulated organic compounds (UOCs) identified in biosolids is also provided (Warke and McAvoy, 2024) (Section 2.4).

## 2.1 PFAS

Per- and polyfluorinated alkyl substances (PFAS) are a group of over 4700 aliphatic compounds containing at least one carbon-fluorine (C-F) bond. Due to environmental concerns, North America, Europe, and Australia voluntarily phased out long-chain PFAS ( $\geq 8$  carbons) in the early 2000s, replacing them with shorter-chain versions. These shorter-chain PFAS are less prone to soil absorption and bioaccumulation, yet they are more environmentally mobile. Despite their reduced bioaccumulation, short-chain PFAS still persist in the environment and can pose risks to human health and the environment (Pozzebon and Siefert, 2023).

Although toxicity and effects data only exist for a small number of PFAS, the health effects of this class of chemicals are well documented. Exposure to PFAS, even at low concentrations, can have significant adverse health effects and effect multiple organs and systems, including liver, kidney and thyroid function, as well as the immune, nervous and reproductive systems (<https://www.canada.ca/en/health-canada/services/environmental-workplace-health/reports-publications/water-quality/water-talk-per-polyfluoroalkyl-substances-drinking-water.html>). Additionally, in the environment, PFAS exposures have been linked to negative impacts on the immune and nervous systems of wildlife, as well as effects on growth, reproduction, and development (ECCC, 2023 and 2024).

In May 2023, Environment and Climate Change Canada and Health Canada published the Draft State of PFAS Report (Government of Canada, 2023), providing a qualitative assessment of the fate, sources, occurrence, and potential impacts of PFAS on the environment and human health. The report highlights the extreme stability of PFAS in the environment, often referred to as "forever chemicals". The report also emphasizes the potential for cumulative effects from co-exposure to multiple PFAS, which may lead to adverse environmental and health outcomes. In July 2024, an updated draft report (Government of Canada, 2024) was issued which incorporated new information and public comments. The updated report continues to underscore the environmental persistence and potential human health and environmental risks associated with PFAS, and reinforces the need for comprehensive management strategies to mitigate their impact. Additionally, in alignment with the Government of Canada's 2021 notice of intent to address PFAS as a class of chemicals (<https://www.canada.ca/en/health-canada/services/chemical-substances/other-chemical-substances-interest/per-polyfluoroalkyl-substances.html>), in August 2024, Health Canada published an objective for Canadian Drinking Water Quality for PFAS of 30 nanograms/L (ng/L) for the sum of 25 PFAS (Health Canada, 2024a).

PFAS have been identified in biosolids globally, and while concentrations of PFAS in biosolids are influenced by inputs to WWTPs, treatment methods, and sludge stabilization techniques, studies have confirmed their presence in biosolids in Canada nationwide (McCarthy, 2015; Letcher et al., 2020; Lakshminarasimman et al., 2021 and Gewurtz et al., 2024).

In response to growing concerns over PFAS in biosolids, in June 2024 the Canadian Food Inspection Agency (CFIA) recommended an interim standard for PFAS in commercial biosolids of 50 part per billion

(ppb) of perfluorooctane sulfonate (PFOS) on a dry weight basis (Canada Food Inspection Agency trade memoranda T-4-132 available at <https://inspection.canada.ca/en/plant-health/fertilizers/trade-memoranda/t-4-132-commercial-biosolids>), with enforcement of the standard to begin as of October 18, 2024. CFIA (2024) indicates that the available data suggest that approximately 92% of Canadian biosolids contain PFOS at concentrations below 50 ppb (ng/g). Data provided by the Capital Regional District (CRD) indicates that the average concentration of PFOS in biosolids from the region is 5.4 ppb (ng/g) and is nearly an order of magnitude below the CFIA limit (see data in Table 1).

The US EPA has prioritized research, restriction, and remediation of PFAS in the environment, including in biosolids, and has defined a PFAS Strategic Roadmap which includes a risk assessment for perfluorooctanoic acid (PFOA) and perfluorooctane sulfonate (PFOS) in biosolids. The assessment is currently underway and is anticipated to be completed by the end of 2024 (<https://www.epa.gov/biosolids/and-polyfluoroalkyl-substances-pfas-biosolids>). During the completion of the assessment, the US EPA has advised states to monitor biosolids for PFAS contamination, identify suspected industrial discharges and implement pre-treatment requirements where appropriate to reduce concentrations of PFAS in biosolids. The US EPA has indicated that if the risk assessment determines that PFOA or PFOS in biosolids may adversely affect public health or the environment, that risk managers will consider options for numerical limitations and best management practices for these compounds.

Additionally in the USA, several states have responded to concerns regarding PFAS in biosolids by enacting rules that limit the concentration of PFAS in biosolids, with Maine (in 2022) and Connecticut (in 2024) implementing bans on the use or sale of PFAS-containing biosolids.

Key findings of recent scientific studies that have evaluated PFAS in biosolids include the following:

1. As the quality of biosolids is dependent on the contaminants present in raw wastewater, it is important to consider upstream controls and monitoring to minimize the input of PFAS-containing wastes into wastewater treatment systems (BC ENV, 2024; Tansel et al., 2024; Gewurtz et al., 2024). In their review of PFAS concentrations in influent, effluent and biosolids samples collected from 27 wastewater treatment plants (WWTPs) across Canada, Gewurtz et al. (2024) identified that PFAS concentrations are the highest in samples from plants that receive landfill leachate.
2. PFAS concentrations in biosolids in the US have not decreased despite the phasing out of the longer chain PFAS in the early 2000s (Pozzebon and Siefert, 2023; Borthakur et al., 2022; Gewurtz et al., 2024). In a study of biosolids sourced from Canadian WWTPs, Gewurtz et al. (2024) reported that except PFOS, the concentrations of long-chain PFAS have generally decreased overtime; however, they remain at measurable concentrations. Further, Gewurtz et al. (2024) found that the replacement of long-chained PFAS with short-chain PFAS has resulted in concentrations of short-chain PFAS in wastewater influent and effluent increasing over the period of 2009 and 2021, with the short-chain PFAS less frequently detected in biosolids.
3. Recent evidence suggests that PFAS are transformed during the wastewater treatment process. Studies (Helmer et al., 2022; Tansel et al., 2024; Behnami et al., 2024) indicate that some PFAS precursors transform during treatments like anaerobic digestion, potentially altering contaminant profiles and increasing certain PFAS concentrations post-treatment. PFAS not degraded during aerobic and anaerobic digestion will instead become concentrated during treatment (Lakshminarasimman et al., 2021; Gewurtz et al., 2024; Tansel et al., 2024). Carbon chain length significantly affects PFAS partitioning, with longer chain PFAS adsorbed to sludge, and shorter chain PFAS partitioning to the wastewater (Behnami et al., 2024; Gewurtz et al., 2024).

## 2.1.1 Screening Level Human Health and Ecological Risk Assessment for PFAS in Biosolids

Data collected from 29 Canadian wastewater treatment plants, as summarized in Zhou et al. (2024) from

Letcher et al. (2020) and Lakshminarasimman et al. (2021) indicate that PFOS was the most prevalent PFAS (<0.49–50.4 ng/g dw) in collected biosolids, followed by PFDA (0.11–53 ng/g dw) and PFOA (<0.07–23 ng/g dw) (Zhou et al., 2024; Letcher et al., 2020; Lakshminarasimman et al., 2021).

Gewurtz et al. (2024) summarized data for 42 PFAS in samples collected in 2021 from 27 WWTP across Canada, and reviewed trends in concentrations overtime. The 2021 results were consistent with the previous studies, indicating PFOS is the most prevalent, with concentrations ranging from <0.32 to 96 ng/g dw.

Gerwurtz et al. (2024) reported that the PFAS with the highest concentrations in Canadian biosolids (2018 to 2021) were PFOS, PFBA and the short-chain precursor 5:3 FTCA. Long-chain PFCAs, PFOS, FOSA, the long-chain intermediate transformation products *N*-methylperfluorooctanesulfonamidoacetic acid (MeFOSAA) and *N*-ethylperfluorooctanesulfonamidoacetic acid (EtFOSAA), PFHxA, and 5:3 FTCA were frequently (in >60 % of the samples) over this period, with the long-chain PFAS likely detected more frequently due to their greater tendency to sorb to solids (Gewurtz et al., 2024; Helmer et al., 2022).

The below table summarizes the concentrations of select PFAS identified in Canadian biosolids (from Letcher et al., 2020; Lakshminarasimman et al., 2021 and Gewurtz et al., 2024). In addition, average PFAS concentrations measured in CRD biosolids (n=3) have been provided for comparative purposes.

PFBA (C4)	PFP eA (C5)	PFHxA (C6)	PFHpA (C7)	PFOA (C8)	PFNA (C9)	PFDA (C10)	PFUnA (C11)	PFDoA (C12)	PFBS (C4)	PFHxS (C6)	PFOS (C8)
Letcher et al., 2020											
<0.48-3.0	<0.28-6.0	0.17-4.65	<0.08-1.53	<0.07-11.5	0.09-4.72	0.11-23.4	0.19-7.49	0.19-6.09	<0.14-3.48	<0.06-2.43	0.49- <b>50.4</b>
Lakshminarasimman et al., 2021											
<2.06	<2.06-14	<2.06-8.3	<2.06-5.2	<2.06-23	<2.06-20	<2.06-53	<2.06-7	<2.06-10	<4.11-11	< <b>4.11</b>	<4.11- <b>27</b>
Gewurtz et al., 2024 (2021 data)											
<1.3-200	<0.64-12	<0.32-7.9	<0.32-2.6	<0.32-42	<0.31-8.8	<0.31-27	<0.31-3.5	<0.31-7.6	<0.32-7.5	<0.32- <b>5.4</b>	<0.32- <b>96</b>
Capital Regional District Biosolids											
<1.50	1.28	2.02	0.96	1.11	0.37	1.80	0.738	1.34	0.474	2.2	5.35
<b>Health Canada SSVs (HC, 2019), BC CSR Standards and CCME Soil Quality Guidelines, Residential/Agricultural Land Use, Human Health Protection</b>											
114,000 <sup>a</sup>	800 <sup>a</sup>	800 <sup>a</sup>	800 <sup>a</sup>	700 <sup>a</sup>	800 <sup>a</sup>	NA	NA	NA	61,000 <sup>a</sup> 300,000 <sup>b</sup>	2,300 <sup>a</sup>	2,100 <sup>a</sup> 1000 <sup>b</sup> 2,000 <sup>c</sup>
<b>Grippo et al., (2021), BC CSR Standards and CCME Soil Quality Guidelines, Residential/Agricultural Land Use, Ecological Protection</b>											
2,980 <sup>d</sup>	NA	6,200 <sup>d</sup>	NA	3,840 <sup>d</sup>	24.2 <sup>d</sup>	67.7 <sup>d</sup>	NA	NA	817 <sup>d</sup>	2.8 <sup>d</sup>	70,000 <sup>b</sup> 10 <sup>c</sup> 8.7 <sup>d</sup>

**Table 1:** Concentrations of PFAS reported in biosolids from WWTPs in Canada (in ng/g) (from Letcher et al., 2020; Lakshminarasimman et al., 2021 and Gewurtz et al., 2024) compared to Health Canada SSVs (2019) and BC CSR soil standards for residential and agricultural land use. NA: Not Applicable, <sup>a</sup> Health Canada SSV, <sup>b</sup> BC CSR soil standard, <sup>c</sup> CCME soil quality guideline, <sup>d</sup> Grippo et al. (2021) ESV, **Bold** – concentration exceeds ecological protection guidelines/standards.

Following the land application of biosolids, there is the potential for human and ecological receptors to be exposed to the biosolids, and thus the contaminants, including PFAS, present in the biosolids. Depending on the land application technique used (e.g., injection, incorporation into surface soils, amendments), the biosolids and COECs may be ‘diluted’. For this screening exercise, it has been conservatively assumed that there is the potential for human and ecological receptors to be exposed directly to the PFAS concentrations measured in Canadian biosolids. In practice, biosolids are mixed with other materials prior to land application; for example the CRD uses a mixing ratio of 18:1 (5 parts sand, 13 parts wood and 1 part biosolids, or approximately 6% biosolids by volume).

Per standard risk assessment methodologies, including those recommended in guidance from the BC ENV

(2023a), Health Canada (2024b) and the CCME (2020), the concentrations of PFAS presented in Table 1 have been compared to guidelines/standards derived to be protective of human health and ecological direct contact exposures, including for human health, incidental ingestion of soil, and for ecological receptors, direct contact, and for wildlife, soil and food ingestion.

Soil guidelines/standards derived to be protective of human health are available for select PFAS from Health Canada (2019), the BC Contaminated Sites Regulation (CSR) (BC, 2023b) and the CCME (2021). Health Canada recommends human health protective soil screening values (SSVs) for 11 select PFAS including PFOS, PFOA, PFBA, PFBS, PFHxS, PFPeA, PFHxA, PFHpA, PFNA and two fluorotelomer sulfonates (6:2 FTS and 8:2 FTS). The SSVs also consider the potential for PFOS and PFOA to act additively, with an SSV of  $< 1$  for the sum of the ratio of the  $[PFOS]/SSV_{PFOS}$  and the  $[PFOA]/SSV_{PFOA}$ . Using the maximum concentration of PFOS and PFOA reported in Table 1 (PFOS = 96 ng/g and PFOA = 42 ng/g), the ratio is  $< 1$  and thus meets the SSV. In addition, the BC CSR provides human health protective soil standards for PFOS and PFBS, and the CCME provides a human health soil quality guideline for PFOS. A comparison of the maximum concentrations of the PFAS measured in Canadian biosolids to the Health Canada SSVs, as well as the BC CSR and CCME human health protective soil standards (See Table 1) indicates that the measured concentrations are well below the SSVs, standards and guidelines, with the average concentrations of PFAS measured in CRD biosolids (n=3) well below the Canadian maximum concentrations and all available SSVs, standards and guidelines. As noted, the direct comparison of measured concentrations in biosolids to the standards and guidelines is highly conservative given the mixing that occurs prior to land application. For example, applying the 18:1 mixing ratio used by the CRD, the concentrations of PFAS in Table 1 would be divided by a factor of 18 to provide a resulting exposure concentration. Applying this ratio, the maximum exposure concentration of PFOS (the PFAS measured at the highest concentration) in CRD biosolids would be 0.3 ng/g compared to the lowest standard for human health protection of 1000 ng/g and the lowest standard for ecological health protection of 8.7 ng/g (protective of food chain exposures).

Soil guidelines and standards in BC and available from the CCME for the protection of the environment, are limited to PFOS only. On behalf of the US Department of Energy, Grippo et al. (2021) developed ecological screening values (ESVs) to support screening-level ecological risk assessments at U.S. Air Force (Air Force), Navy, Army, and other U.S. Department of Defense (DOD) sites PFAS have been detected in soils and surface waters. A comparison of the maximum concentrations of PFAS measured in Canadian biosolids to the ESVs and CSR/CCME guidelines/standards for PFOS is provided in Table 1. As presented, concentrations of select PFAS, including PFOS and PFHxS exceed the guidelines for environmental protection, and specifically the CCME soil quality guideline and the Grippo et al. (2021) EVS for soil and food ingestion. Concerning the Grippo et al. (2021) ESV that is exceeded, it is specific to soil and food ingestion of mammalian ground insectivores. Further, concentrations of PFOS exceed the CCME guidelines protective of soil leaching to groundwater for the protection of potable groundwater (10 ng/g) and aquatic life (10 ng/g).

The below discussion considers screening quotients (SQs) for select PFAS; SQs were calculated as the  $[PFAS] / \text{guideline (or standard or screening value)}$ .

The above screening exercise indicates that the concentrations of individual PFAS in both Canadian biosolids, including those from the CRD, for which there is available toxicity data to derive soil guidelines are less than the guidelines derived to be protective of human health (maximum SQ of 0.06 for PFOA), and thus, exposure to these individual PFAS in biosolids, as well as combined exposures to PFOS and PFOA, are not anticipated be associated with risks to human health. As noted the comparison of measured concentrations is highly conservative as amendment would occur prior to application.

The comparison of the PFAS concentrations measured in Canadian biosolids, as reported by Letcher et al., 2020; Lakshminarasimman et al., 2021 and Gewurtz et al., 2024 exceed ecological health guidelines for select individual PFAS including PFOS (maximum SQ = 11) and PFHxS (maximum SQ = 1.9). On this basis, there is the potential for the highest concentrations of these PFAS measured in Canadian biosolids



to pose a risk to ecological receptors and specifically wildlife exposed via soil and food ingestion. Further, as the maximum PFOS concentrations exceed of soil guidelines protective of drinking water and aquatic life (maximum screening quotient of 9.6), there are potential for risks to both human health and the environment associated with soil leaching to groundwater. The average concentrations of PFAS in the CRD biosolids were all below the available SSVs, ESVs, as well as the CCME guideline and CSR standard for PFOS (i.e., all SQs are well below 1).

The findings of the above screening exercise must be interpreted in the context that guidelines are only available for a small number of the 4700 PFAS known to exist, and that the individual guidelines may not consider the combined toxicity of the PFAS mixture present in biosolids. Further, as PFAS are persistent and accumulate in soils, following the repeated application of biosolids to land, the concentrations of PFAS have the potential to increase overtime.

## 2.1.2 Estimated Daily Intakes of PFOS and PFOA

As PFAS are ubiquitous in the environment, humans are exposed daily to PFAS from a variety of sources (i.e., consumer products, diet, air, water and soil). Limited data exists on Canadian's exposure to PFAS; however, biomonitoring data indicates that PFAS are present at measurable concentrations in the blood of most Canadians (Government of Canada, 2024).

Using summary statistics from secondary sources for concentrations of PFOA and PFOS in indoor and outdoor air, water and dust in the US, as well as European dietary intake estimates to estimate exposures from food, East et al. (2023) estimated exposure to adults and children over the period of 2011 to 2017. Daily intake estimates for adults were estimated to be 40 ng/day PFOA and 40 ng/day PFOS, and rates for young children (toddlers) were estimated to be 14 ng/day PFOA and 17 ng/day PFOS. A comparison of these estimates using a first-order pharmacokinetic model indicated that the results were aligned with serum concentration measurements from the National Health and Nutrition Examination Survey over the same time period (East et al., 2023), providing evidence that the modeled daily intakes are reasonable. It is noted that although not discussed by East et al. (2023), it is assumed that exposures to PFAS in biosolids would be accounted for in the measured serum contributions in areas where biosolids are applied and represent a potential source.

As measured levels of PFAS in blood in Canada and the US are reportedly similar (Public Health Ontario, 2023), the results of East et al. (2023) have been considered here in an assessment of potential PFAS exposures for Canadians, and a comparison to potential exposures to PFOS and PFOA measured in Canadian biosolids.

Using Health Canada (2024b) exposure equations for the direct soil exposure pathways (i.e., incidental ingestion, dermal contact and inhalation of soil particulate) and receptor characteristics for an adult and a toddler, with the average PFOS and PFOA concentrations in CRD biosolids from Table 1 used as exposure point concentrations, exposure intakes associated with exposures to PFOA and PFOS in CRD biosolids were estimated. A summary of the results summed with the estimates from East et al. (2023) and compared to the Health Canada tolerable daily intakes (TDIs) for PFOA and PFOS are presented in Table 2.

PFAS	Background EDI (East et al., 2023) (ng/day)		EDI from CRD biosolids (ng/day)		Total Exposure Estimate (ng/day)		Total Exposure Estimate (ng/kg-bw/day)		Health Canada Tolerable Daily Intakes (ng/kg-bw-day)
	Toddler	Adult	Toddler	Adult	Toddler	Adult	Toddler	Adult	
PFOA	14	40	0.09	0.03	14.09	40.03	0.85	0.57	60
PFOS	17	40	0.44	0.14	17.44	40.14	1.06	0.57	21

Table 2. Estimated daily intakes for PFOA and PFOS (from East et al., 2023) and from exposures to PFOA and PFOS in biosolids using the maximum PFOA and PFOS concentrations in Table 1. Total exposure estimates are the sum of the EDIs from East et al., 2023 and the EDIs from biosolids.

The total exposure estimates (the sum of the estimated daily intakes from East et al., 2023 and those from CRD biosolids) in ng/kg-bw-day are well below the Health Canada (2021) tolerable daily intakes (TDIs),

suggesting that exposures for PFOA and PFOS in biosolids in the CRD, combined with background exposures, are unlikely to represent a health risk. The estimated exposures from the CRD biosolids are conservative as they are based on concentrations measured in the CRD biosolids, which as noted, would be reduced by a factor of 18 following the amendment of the biosolids prior to application.

As with the results of the screening assessment presented in Section 2.1.1, the above evaluation does not consider exposures to the mixture of PFAS in biosolids, which has the potential to act additively with one another as well as with other contaminants in biosolids, and thus is likely an underestimate of total risks associated with PFAS exposures. Despite this, the potential contribution of exposures to PFOS and PFOA from biosolids from the CRD to overall exposures of these PFAS is low, and when amendment of the biosolids is considered, is likely to be negligible.

## 2.2 Microplastics

Microplastics are pieces of plastic less than five millimeters in diameter and while some microplastics are intentionally manufactured (e.g., microbeads in beauty products), they generally result from the degradation of plastic debris. Microplastics have been identified to be present in biosolids around the globe, including in Canada (Crossman et al., 2020; Gies et al., 2018; Lavoy and Crossman, 2021; Sivarajah et al.; 2023). Further, biosolids have been identified as an important pathway for microplastics to enter the environment (Crossman et al., 2020) and agricultural lands where biosolids have been applied are one of the largest reservoirs of microplastics (Pozzebon and Siefert, 2023). Due to their emerging nature, there are no regulations pertaining to microplastics in biosolids (or from other sources) in Canada.

While the studies assessing the prevalence of microplastic in biosolids in Canada were previously limited, Sivarajah et al., (2023) quantified microplastics in biosolids from 22 WWTPs located in nine Canadian provinces. Microplastics were identified in all samples, at concentrations ranging from 228 to 1353 particles per gram dry weight (dw) (median = 636 particles per gram dw). These concentrations are orders of magnitude greater than those reported from previous investigations of microplastics in biosolids in 4 other WWTP in Canada, as well as in other countries (Sivarajah et al., 2023). Importantly, despite the large variation in the concentrations of microplastics observed across the samples from the 22 WWTPs, the investigators did not find a significant difference in the concentrations based on region, the type of WWTP or the sludge treatment type. The results for the Pacific region (n=4) were the highest, with a median concentration of 914 particles per gram dw.

The effects of microplastics on humans and the environment remain largely uninvestigated. In recognition of this, in January 2024 the Government of Canada announced funding of \$2.1 million over four years to three academic institutions for the research of microplastics and their potential to impact human health (<https://www.canada.ca/en/health-canada/news/2024/01/government-of-canada-funding-research-on-the-health-risks-of-microplastics.html>). While there is limited data on health effects, the inhalation of microplastics has been identified as a concern, with the inhalation of microplastics associated with oxidative stress in lung tissues and general inflammation responses in airways (Pozzebon and Siefert, 2023).

Microplastics are highly persistent, resistant to degradation and can accumulate in soils (Xu et al., 2019; Pozzebon and Siefert, 2023). This is of specific concern in areas where there is repeated land application of biosolids. Contrary to the benefits of the land application of biosolids, microplastics from biosolids and their accumulation in soil compromises soil structure and affects nutrient availability, water retention and aeration (Xu et al., 2019). They can also be toxic to soil organisms and thus reduce the beneficial effects these organisms have on soil fertility and structure (Xu et al., 2019). Evidence also suggests the potential for the chemical constituents in microplastics to leach into soil and groundwater, and for plants to absorb microplastic particles, serving as a pathway for microplastics to enter the food chain (Xu et al., 2019; Pozzebon and Siefert, 2023).

A recent study (Wang and Good, 2024) identified that microplastics in biosolids can act as vectors for the long-range transport of PFAS, including atmospheric deposition in aquatic systems. The authors indicate

that microplastics enriched with PFAS are an important concern due to the ubiquitous nature of both microplastics and PFAS globally, but also their co-occurrence in biosolids and the potential for combined toxicity (Wang and Good, 2024).

The presence of microplastics in biosolids, and the early evidence of effects to both human health and the environment, highlights the need for further research and the implementation of measures to prevent the further introduction of microplastics to the environment. Recent policies limiting single use plastics, along with existing regulations banning use of polymeric microbeads in cosmetics and personal care products, are likely to have resulted in a reduction in microplastics entering WWTPs. Despite this, the laundry of synthetic clothing will continue to contribute to the load of microplastics in biosolids (Crossman et al., 2020); however, the use of microfibre filters in washing machines would reduce microplastics sourced from laundering clothing.

## 2.3 Pharmaceuticals and personal care products

Pharmaceutical and personal care products (PPCPs) include prescription and over the counter medications and supplements, as well as pharmaceuticals used in agriculture to promote growth and health of livestock (Pozzebon and Siefert, 2023; Hydromantis Inc., 2010; McCarthy, 2015). PPCPs also include other products used for health and cosmetic purposes (e.g., fragrances). Sub-categories of PPCPs include antibiotics, antimicrobials, steroidal chemicals, fragrances, alkylphenolics, and other PPCPs such as analgesics, antidepressants, antifungals, anti-inflammatories and diuretics, to name a few.

Traditional WWTPs are generally ineffective at removing PPCPs, which often persist through the treatment process. The land application of biosolids has been identified as the primary way that PPCPs enter the environment (Brown et al., 2019; Kinney and Vanden Heuvel, 2020; Pozzebon and Siefert, 2023).

Data indicate that most PPCPs degrade within months after being introduced to the environment; however, some studies indicate that select PPCPs can persist (Garcia-Santiago et al., 2016; McCarthy, 2015; Kinney and Vanden Heuvel, 2020). Further, many PPCPs are considered "pseudo-persistent" as they are continuously introduced to the environment (Pozzebon and Siefert, 2023). The potential for PPCPs to accumulate in soil and transfer to soil invertebrates and plants, as well as to leach to groundwater has been demonstrated (Garcia-Santiago, 2016), although limited data exists on the ecotoxicity of PPCPs, especially in terrestrial systems (Pozzebon and Siefert, 2023). Given the low doses of many PPCPs required to elicit a response in humans, PPCPs are likely to cause a variety of effects on biota if they accumulate in the environment. PPCPs contain endocrine disruptors which have the potential to interfere with human and animal hormonal systems even at low concentrations. Reproductive failure associated with exposure to endocrine disrupting contaminants has been well documented in aquatic systems (Pozzebon and Siefert, 2023).

Further, some antibiotics persist in the environment after land application, affecting soil microbial communities and potentially altering ecosystem functions. They can promote the growth of antibiotic-resistant bacteria, and may disrupt the balance of beneficial soil microorganisms, affecting nutrient cycling and soil health (Black et al., 2019; Pozzebon and Siefert, 2023). Antibiotic resistance has been defined by the World Health Organization as a global threat to health and food security. The promotion of antibiotic-resistant bacteria by PPCPs will contribute to this threat and is specifically concerning given the co-occurrence of antibiotic-resistant bacteria and pathogens in biosolids (Pozzebon and Siefert, 2023). Hung et al. (2022) found that biosolid samples contained significantly higher concentrations of antibiotic-resistant genes when compared to raw agricultural soils, as well as the potential for the airborne spread of the genes. The authors emphasized the importance of their findings to the global concerns regarding antibiotic resistance.

Previous studies (Garcia-Santiago et al., 2016; Kennedy/Jenks, 2017) have conducted preliminary evaluations of the potential effects and risks associated with PPCPs. Additionally, McCarthy (2015) summarized the results of previous risk assessments including those for PPCPs. Overall, the assessments



suggest that the potential for human health risks from exposure to individual PPCPs in biosolids is likely low, however, there are limitations and uncertainties in the previous assessments, including the consideration of a small number of PPCPs and the lack of accounting for potential additive and synergistic effects.

Garcia-Santiago et al. (2016) conducted a screening level risk assessment to assess the potential for human exposure to PPCPs measured in biosolids to exceed one therapeutic dose. The study focused on PPCPs shown to persist in WWTPs including carbamazepine, fluoxetine, triclosan, miconazole and ciprofloxacin, and Naproxen. Total exposures via the direct soil exposure pathways, as well dietary exposures, were evaluated. The results of the assessment indicated total hazard quotients (HQs) were less than an HQ of 1.0 (i.e., total exposures were less than one therapeutic dose), with most PCPPs having HQ values of less than 0.1, except for triclosan which has been shown to bioaccumulate, which had an average HQ of 0.28 and a 95% UCLM HQ of 0.95, indicating a potential risk to human health. The mean triclosan concentration reported by Garcia-Santiago et al. (2016) was 5,890 (SD = 3,870) ng/g, while the concentration of triclosan measured in CRD biosolids (n=1) is 1,870 ng/g.

Kennedy/Jenks Consultants (Kennedy/Jenks, 2017) completed a biosolids risk assessment on behalf of Metro Vancouver. A quantitative human health risk assessment was conducted for a small group of COECs detected in biosolids including over the counter pain medications (analgesics) and non-steroidal anti-inflammatory drugs, antidepressants, antibiotics, antimicrobials, plasticizers, and flame retardants. The results of the risk assessment suggested that the concentrations of the COECs in biosolids from the Metro Vancouver region were unlikely to result in adverse health effects to exposed individuals, including children. The assessment further demonstrated that it would take a minimum of a decade and up to one billion years of exposures to the COECs in biosolids to equal a single therapeutic dose of the PPCPs.

The number of PPCPs identified in biosolids continues to grow and there is a general lack of data on the effects of these chemicals in the environment. Given that many of them act via the same mode of action (e.g., endocrine disruptors) and/or belong to the same classes (e.g., antibiotics, SSRIs) it is essential that future assessments consider the combined effect of the mixture of these PPCPs on human health and the environment.

## 2.4 Prioritization of Unregulated Organic Chemicals in Biosolids (Warke and McAvoy, 2024)

While acknowledging the benefits of the land application of biosolids, Warke and McAvoy (2024) conducted a literature review to compile a database of all reported unregulated organic chemicals (UOCs) present in biosolids. Where data gaps were identified, predictive modelling and an extensive literature search were conducted to determine values for persistence, mobility, bioaccumulation, and toxicity. The prioritization process used these characteristics to rank the UOCs according to their potential impact on human health.

Of 906 chemicals identified in biosolids, 124 were categorized as either high or low priority. Among these, 13 chemicals were classified as carcinogenic, and 22 as endocrine disruptors. Notable examples of endocrine-disrupting chemicals included N-nitrosodimethylamine, cashmeran, nonylphenol, bisphenol A, and several PBDEs. Potential carcinogens identified included 1,2-dichloropropane, 1,4-dioxane, di(2-ethylhexyl)-phthalate, and trichloroethylene.

The priority UOCs were further ranked using scoring based on combinations of mobility, persistence, bioaccumulation, and toxicity. This analysis added eight additional compounds to the high-priority list, resulting in a total of 46 high-priority compounds, with the remaining 78 classified as low priority. The high priority UOCs included several carcinogens (e.g., N-nitrosodiethylamine), endocrine disruptors (e.g., BDE 99, estrone), PPCPs (e.g., fluoxetine, bisphenol-A, carbamazepine, triclosan) and industrial solvents (e.g., trichloroethylene)

Comparison of the Warke and McAvoy (2024) results to those from other studies ranking UOCs in biosolids identified similarities. Of the 46 high-priority UOCs identified by Warke and McAvoy (2024), 38 were also present in other priority lists, including 20 in from Higgins et al. (2010), 14 in a study conducted on behalf of Scottish EPA (WCA, 2019), and 12 in a study conducted by the Texas Commission on Environmental Quality (TCEQ, 2021).

Of the high-priority UOCs identified, several including: bisphenol-A, triclosan, nonylphenol, N-nitrosodimethylamine, 4-chloraniline, triphenyltin (TPhT), and several polybrominated diphenyl ethers (PBDEs) such as BDE 209, BDE 47, and BDE 99, have been associated with effects ranging from endocrine disruption to carcinogenic effects (Warke and McAvoy, 2024). Additionally, triclosan, is known to disrupt thyroid function, may cause reproductive and developmental toxicity and N- nitrosodimethylamine has been linked to neurological, gastrointestinal, and developmental disorders (Warke and McAvoy, 2024).

As noted, previous risk assessments for biosolids-amended soil indicate that most UOCs are below threshold levels for human exposure pathways ( $HQ < 1$ ). However, Warke and McAvoy (2024) emphasized that preliminary data for pathways involving soil organisms and aquatic systems suggest the potential for HQ values exceeding 1 for certain UOCs including triclocarban, ciprofloxacin, and azithromycin, and that further study is needed for these pathways. Further, while some compounds like caffeine pose minimal risks to humans, they can significantly impact aquatic and soil organisms and antibiotics like ciprofloxacin and azithromycin, as well as fluoroquinolones, have been identified as photosynthetic inhibitors in plants and present potential risks to soil and aquatic ecosystems (Warke and McAvoy, 2024).

While the available data generally support that the measured concentrations of PCPPs in biosolids are low and for the most part do not represent a risk to human health and the environment, the results of Warke and McAvoy (2024) further highlight the number of contaminants in biosolids, and gaps where further research is required to understand potential risks to human health and the environment.

### 3 Fate and Transport of Contaminants in Biosolids and Considerations of Conditions Typical of the BC South Coast

In sensitive coastal environments, like those present in southern BC, the fate and transport of contaminants, is complex. As discussed throughout this report, land application of biosolids can result in the introduction of a range of contaminants, which may disperse in the environment through soil leaching, erosion, and runoff. This issue is exacerbated by heavy precipitation which can enhance the mobilization of contaminants. Precipitation infiltrates the soil and can leach soluble contaminants into groundwater. In agricultural and rural areas, where biosolids are most likely to be applied, the use of groundwater as a source of drinking water, irrigation water or livestock water, is more likely. Thus, the contamination of groundwater has the potential to result in human health exposures via potable water, as well as livestock and crop exposure. Further, groundwater impacted with contaminants from biosolids can migrate to nearby aquatic systems and result in effects to aquatic ecosystems, as well as entry into the human food chain via seafood consumption.

Heavy rainfall or snowmelt can also create surface runoff, which has the potential to mobilize contaminants from application areas, resulting in impacts to adjacent lands, or migration to nearby water bodies. For example, Crossman et al. (2020) suggested that heavy rainfall may result in the mobilization of microplastics from agricultural soils. The risk of contaminant mobilization via surface runoff is especially high prior to the incorporation of applied biosolids into the soil matrix (LRCS Land Resource Consulting Services, 2016).

Further, prolonged or intense precipitation can saturate soils, resulting in decreased soil adsorption of contaminants and subsequent leaching, as well as soil erosion. Eroded soils carrying adsorbed contaminants may be redistributed across the landscape onto adjacent lands or to nearby waterbodies. Additionally, natural processes such as freeze-thaw and drying cycles can release fine particles from biosolids, facilitating the movement of contaminants into subsurface soils and groundwater.

Tansel et al. (2024) discussed that PFAS persistence and mobility are dependent on soil interactions and precipitation events. Leaching, which is influenced by soil type and water infiltration rates, with higher infiltration rates in areas with high precipitation levels, facilitates PFAS migration to groundwater and surface water. PFAS compounds with long fluoroalkyl chains tend to bind more strongly to solid phases, while shorter-chain PFAS are soluble and prone to leaching and transport. Consistent with other studies (Blake and Fenton, 2020; Drew et al., 2021), Tansel et al. (2024) identified the potential for plant uptake of PFAS from biosolid-amended soils, raising concerns about bioaccumulation in food chains and ecosystems.

PCPPs in biosolids can similarly migrate, with pharmaceuticals showing potential to leach into groundwater (Santiago et al., 2016; Kinney et al., 2006). Gottschall et al., (2012) detected pharmaceuticals and personal care products (PPCPs) in groundwater following biosolids land application. While many PPCPs dissipate within a few months, others, particularly those embedded in biosolid aggregates, can persist for over a year (Kinney and Heuvel, 2020).

Some COECs, due to their persistence and bioaccumulation potential, can be transported long distances. For example, microplastics in biosolids exhibit hydrophobic characteristics, allowing them to adsorb other contaminants. These particles may then be subject to long-range transport (Carbery et al., 2018). Studies (Wang et al., 2024, Strynar et al., 2011) indicate that long-chain PFAS can attach to microplastics or dust and become airborne. These findings raise concerns regarding the multiple contaminant types present in

biosolids and their influence not only on the fate and transport of such contaminants, but also on their combined toxicities.

While the science on the fate and transport of COECs in biosolids is advancing, further research is required to fully understand the behaviour of these contaminants in the environment. Field studies in BC that evaluate seasonal influences, such as precipitation levels, on the environmental fate and transport of COECs in biosolids would provide valuable insight, including how the co-occurrence of numerous contaminants impact migration patterns and ultimately exposure pathways for human and ecological receptors.

## 4 Techniques to Reduce Risks in Biosolids Land Application

Numerous studies have emphasized the importance of government oversight and regulation to limit the risks associated with the land application of biosolids; however, as discussed, available regulations are limited due to the paucity of toxicity data for COECs. This gap, and the uncertainties discussed in Section 6, highlight the critical need for further research, source control to limit the entry of COECs into WWTPs, the development of treatment technologies that degrade COECs in biosolids, and standardized requirements for monitoring COECs in biosolids.

### 4.1 Recent Research on Treatment Technologies

Recognizing the critical problem that contaminants, and specifically COECs, in biosolids present, recent research has focused on treatment technologies that remove COECs. As noted, conventional WWTPs were not designed to remove COECs and thus, these contaminants will continue to persist in inputs to WWTPs.

Much recent research has been focused on technologies that destroy PFAS. An PFAS Innovative Treatment Team was formed by the USEPA in 2020 to investigate and develop innovative tools and methods to break the carbon fluorine (C-F) bonds in PFAS-containing waste. Four emerging technologies were identified by the team with a technology's suitability dependent on waste characteristics, processing requirements, and potential byproducts, as well as considerations for energy consumption, costs, and system mobility (Berg et al., 2022). Berg et al. (2022) also indicated that additional pretreatment and post-treatment steps may be necessary to enhance effectiveness and manage byproducts, such as volatile PFAS emissions.

The four emerging technologies were summarized by Berg et al. (2022) and include:

- Mechanochemical destruction (MCD), which has been shown to result in over 99% PFAS destruction in laboratory settings. The authors note that this technology requires further study for commercial-scale application.
- Electrochemical oxidation (EO), which uses electrical currents to break C-F bonds in PFAS, has shown successful bench and pilot-scale results, but faces challenges in scaling up.
- Supercritical water oxidation (SCWO) treats waste at high temperatures and pressures, producing heat that can sustain the process, but requires managing acidic byproducts and salt precipitation.
- Pyrolysis and gasification decompose materials at high temperatures, potentially destroying PFAS, and produce useful byproducts like char and syngas. Further research on this technology was noted to be required.

Research briefs on these technologies are available on the PITT's website, providing detailed information on benefits and areas needing further research available at <https://www.epa.gov/chemical-research/pfas-innovative-treatment-team-pitt>. Each of these technologies were noted by US EPA to be under evaluation, with further pilot-testing and reporting of the results planned for 2021; however, no further updates on these technologies were identified on the above webpage. While showing promise, the technologies are not yet commercially available.

Keller et al. (2024) pyrolyzed biosolid samples from a WWTP in Southern California at temperatures ranging from 400 to 700 °C for two hours. The study evaluated contaminant removal, with most contaminants being eliminated entirely and only minimal residuals detected. Notably, no PFAS were detectable at the lowest temperature tested (400 °C), and overall removal of PPCPs exceeded 99.9%. Microplastic removal ranged from 91 to 97% depending on conditions. Additionally, the resulting biochar was rich in iron and phosphorus,

making it a valuable fertilizer additive. The authors of the study indicated that a techno-economic analysis showed that pyrolysis could lead to significant cost savings, with revenue from biochar sales having the potential to offset the capital and operational costs of the drying and pyrolysis systems.

Vo et al. (2024) assessed treatment technologies for microplastics and organic contaminants in biosolids. Their multi-criteria analysis identified anaerobic digestion as the most established and practical approach, indicating that while thermal treatment shows potential, the application requires further advancements in infrastructure, regulatory frameworks, and public acceptance to become widely viable.

Recent studies have explored advanced oxidation treatment methods which are reportedly effective at degrading COECs including PPCPs and PFAS. Booton et al. (2024) indicate that chemical oxidation offers a promising alternative and eliminates persistent contaminants. Compared to biological systems, it is potentially simpler to operate and maintain, while requiring less space for efficient treatment. Key advantages of chemical oxidation are reported by the authors to include its ability to address COECs, its rapid start-up and shutdown capabilities, and its ability to avoid common challenges in biological treatment, such as toxic load management, biomass washout, sludge settling issues, and the complexities of sludge handling and disposal.

The recent research illustrates the potential for available technologies to destroy COECs, demonstrating that with further testing and implementation, that the risks associated with the land application of biosolids could be greatly reduced.

## 4.2 Application Techniques and Site-Specific Application Considerations

Specific application techniques, such as injection, surface incorporation and amendment/mixing of biosolids to produce a biosolids growing media, have been demonstrated to reduce risks associated with biosolids land application. Injecting biosolids below the soil surface reduces bioaerosol dispersion, wind and water erosion, and prevents exposure to contaminants by reducing dust generation and the potential for surface contact. Injection also reduces adherence to plant tissues, and therefore exposures to livestock and wildlife. Similar results may be achieved by mixing biosolids into soil immediately after application (LRCS Land Resource Consulting Services, 2016).

BC ENV (2008), LRCS Land Resource Consulting Services (2016) and others have provided recommendations for reducing risks associated with the land application of biosolids. The recommendations include risk-based planning to reduce exposures, including avoiding sensitive ecosystems and proximity to water bodies, as well as areas with shallow groundwater where leaching of contaminants is more likely. Further, the use of personal protective equipment for workers, adhering to buffer zones required in the OMRR and avoiding application during heavy rainfall and snowmelt will further reduce exposures and the potential for contamination of groundwater and surface water. Implementing waiting periods between application and livestock (and wildlife) exposures is also recommended.

The amendment of biosolids with biochar or wood chips, as done in the CRD, has also been demonstrated to enhance the degradation and/or retention of leachable PPCPs (Pozzebon and Siefert, 2023).

## 5 Limitations of Extrapolating Lab-Based Testing to the Environment

The limitations in extrapolating lab-based toxicity testing results to real-world environmental scenarios are generally recognized and have been well documented by others (Cairns, 1983; Smith and Cairns, 1993; Hill et al., 1994). These limitations are exacerbated when considering the land application of biosolids owing to the large number of contaminants, including COECs, present in biosolids, and as the fate and toxicity of these contaminants which will vary depending on the mixture present and the application site characteristics. Controlled laboratory conditions cannot replicate the complex, variable nature of these environments.

The key limitations that arise during this extrapolation process include:

1. Laboratory tests are conducted under highly controlled conditions, which typically include simplified systems that do not reflect the complexity of the natural environment. Factors such as contaminant mixtures, environmental conditions (e.g., soil type, pH), and contaminant exposure pathways cannot be replicated in a lab setting.
2. Lab tests typically assess the effects of a single chemical or a small group of related chemicals. In the environment, specifically in the case of biosolids, organisms are exposed to complex mixtures with potential additive, synergistic, or antagonistic effects.
3. Lab studies typically expose test organisms to high concentrations of a single contaminant over short durations in confined spaces, with even chronic tests typically limited in duration. This does not reflect environmental exposures which often involve chronic exposures to low levels of contaminants over large areas. The effects of chronic, low dose exposures are typically underestimated in lab settings, leading to potential inaccuracies when predicting chronic toxicity in the natural environment.
4. In toxicity testing, exposures are usually simplified and typically limited to immersion in a contaminated medium or direct ingestion or inhalation. In natural environments contaminants exposure pathways are more complex and may include cross-media exposures, including food chain exposures. This oversimplification has the potential to underestimate exposures and associated effects.
5. Lab toxicity test methods have been developed for a limited number of species, focusing on model organisms that may not be representative species that are most sensitive to a specific contaminant. Further, lab tests may not include life stages (e.g., juveniles, larvae) most sensitive to the contaminants tested.
6. Laboratory organisms are often maintained in stable environments and lack the physiological adaptations that organisms may develop in response to natural stressors. Additionally, organisms may behave differently in the lab setting compared to in their natural environment, potentially influencing their exposure and response to contaminants.
7. In natural environments, interactions with abiotic factors (e.g., soil composition) and biotic factors (e.g., predation) may influence toxicity. These factors cannot be accounted for in a laboratory.

The above limitations of laboratory-based toxicity studies highlight the need for caution when extrapolating laboratory testing results to the environment. As noted, given the complex contaminant mixtures known to be present in biosolids, the likely potential for synergistic or additive effects, as well as the influence of the characteristics of the application area on fate and toxicity, the importance of field studies in the assessment

of potential risks from the land application of biosolids cannot be understated. Field observations, together with laboratory toxicity testing, are essential to understanding risks. The collective results of both, once available, should be considered in the establishment of a risk-based, adaptive management strategy for the land application of biosolids.



## 6 Uncertainties in the Risks Associated with the Land Application of Biosolids

Numerous uncertainties exist in the assessment of risks associated with the land application of biosolids. Many of these uncertainties have been documented by others (McCarthy, 2015; Pozzebon and Siefert, 2023; Garcia-Santiago et al., 2020; Schoof and Houkal, 2005; LRCS, 2016), including those summarized below and discussed in previous sections of this report. The rapidly evolving science on COECs in biosolids and their fate and effects following land application, as well as the very recent government policies and regulations pertaining to COECs in biosolids in Canada and elsewhere (e.g., CFIA October 2024 limit for PFAS in biosolids; Government of Canada July 2024 draft report of PFAS; Government of Canada 2023 funding for research on microplastics), underscore that the science on COECs is “emerging”.

Some of the key scientific gaps and uncertainties related to the land application of biosolids are summarized below:

1. Regulatory agencies have not derived limits for most COECs in biosolids, and further, given the paucity of toxicity (including ecotoxicity and specifically for wildlife at all trophic levels) and fate data for COECs, have not derived environmental quality standards. For the same reason, risk assessments evaluating the potential human health and environmental risks associated with exposures to COECs in biosolids, are limited and have generally focused on only a few of the more common COECs.
2. Due to advances in analytical chemistry methodologies, new COECs present in biosolids continue to be identified. The fate and effects of these contaminants are not well understood and thus, it is not possible to assess the potential for risks to human health and the environment.
3. Existing risk assessments and other evaluations of COECs in biosolids are focused on individual contaminants and are based on toxicity data from laboratory toxicity tests. These risk assessments do not consider that exposures to biosolids would result in the simultaneous exposure to numerous contaminants, and the potential for the contaminants to act additively or synergistically and thus are likely to have underestimated the potential for risks.
4. Similarly, the coexistence of numerous contaminants in biosolids may affect their fate, transport and distribution and therefore potential exposure pathways for human and ecological receptors. As an example, Wang and Good, (2024) identified that microplastics in biosolids can act as vectors for the long-range transport of PFAS. Field studies evaluated the influence of the contaminant mixtures present in biosolids on their environmental fate and transport are required to address this uncertainty.
5. The limited available toxicity data for most COECs is based on laboratory toxicity testing. The limitations and uncertainties associated with lab studies, and the need for field studies to be considered in the interpretation of risks, is highlighted in Section 5.
6. The long-term impacts of COECs in biosolids after land application, including their potential to accumulate overtime, leach into groundwater, enter the food chain, or impact human health or wildlife, are not fully understood. Further research in these areas is essential for the assessment and management of risks.
7. Efficient technologies for detecting and measuring COECs in biosolids and the environment are underdeveloped. Without reliable monitoring, it is challenging to assess or mitigate potential risks associated with contaminants in biosolids.

The rate at which our understanding of COECs in biosolids is advancing, with new data on the fate and effects of COECs being published continuously, underscores the importance an adaptive management framework for the land application of biosolids. . With time, it is anticipated that the uncertainties and data gaps identified here and elsewhere will be addressed, and thus regulators must keep pace with the evolving science and regularly weigh the risks and benefits of the practice in an informed and transparent manner.

## 7 Biosolid Land Application Bans

Several countries have restricted or banned the land application of biosolids due to concerns about environmental contamination, health risks, and public opposition primarily related to the presence of PFAS in biosolids, and their associated entry into the environment, specifically the human food chain.

In Canada, the land application of biosolids has been banned in specific regions due to environmental and public health concerns. In British Columbia, the Capital Regional District implemented a ban on the land application of biosolids in 2011. The ban was implemented due to concerns regarding COECs in biosolids, and to protect local drinking water sources, the environment, and public health. In March 2023, Quebec announced a temporary ban on the import and land application of biosolids originating from the United States due to concerns over PFAS contamination in imported biosolids. The province is working towards establishing standards for PFAS in biosolids to ensure environmental safety.

The following is a summary of the various countries that have banned or implemented strict regulations on the land application of biosolids.

### 7.1 United States of America

In the U.S.A., some states and municipalities have imposed restrictions or moratoriums on the land application of biosolids. In 2022, Maine became the first state to ban the land application of biosolids after it found PFAS had contaminated crops or water on over 50 farms throughout the state where sludge had been spread (Carey, 2023). Other states are starting to implement limits and bans. As of October 1, 2024, Connecticut prohibited the sale of PFAS-containing biosolids or wastewater sludge. Further, Michigan, New York, and Wisconsin have implemented interim strategies that limit the PFAS concentrations allowed in land-applied biosolids, and Colorado's interim strategy requires Source Control Programs to evaluate potential PFAS sources if concentrations in biosolids exceed a determined level.

Several more states have pending legislation that would enforce similar restrictions. As an example, Massachusetts is developing legislation that would set maximum levels for the amount of PFAS allowed in any fertilizer sold in the retail market, and proposed legislation in Oklahoma would require a warning label on any product derived from biosolids or sewage sludge.

### 7.2 Europe

In Europe, several countries have adopted a precautionary approach in banning the land application of biosolids, emphasizing soil and food safety. Switzerland has had a ban on agricultural use in place since 2006 with their current regulation requiring sewage sludge to be combusted, while the Netherlands has imposed stringent limits on several contaminants commonly identified in biosolids, resulting in very limited use. Due to concerns over COECs, including PFAS, pharmaceuticals and microplastics, Sweden, Germany and Austria (Vienna region) have banned the application of biosolids on agricultural land. The ban in Germany was implemented with the amendment of the German Sewage Sludge Ordinance in 2017, and by 2029 biosolids land application will be phased out. Germany is reportedly shifting towards incineration of sewage sludge and phosphorus recovery.

## 8 Discussion and Conclusion

In British Columbia, approximately 38,000 tonnes of biosolids are produced annually, with approximately 72% of biosolids and biosolids-derived products applied to land (BC ENV, 2019). As discussed throughout, while there are many benefits associated with the land application of biosolids, biosolids contain a complex mixture of contaminants, including COECs. Data on the fate and effects of these COECs is limited, and our understanding of the risks that these contaminants present to human health and the environment is rapidly evolving.

When weighed collectively, the available information on the land application of biosolids, including the information presented by others (e.g., Pozzebon and Siefert, 2023; LRCS Land Resource Consulting Service, 2016; McCarthy, 2015; BC ENV, 2024) and herein, supports the assessment of the practice in the context of the Precautionary Principle. The Precautionary Principle guides decision-makers to take action to protect the environment and public health in the face of environmental or health uncertainties (Goldstein, 2001).

Given the significant data gaps and uncertainties in the land application of biosolids, as well as the rapidly advancing science, previous studies that have concluded a low risk to human health and the environment must be interpreted in the context of the uncertainties. Risk assessments are conducted using the best scientific evidence available at the time of the assessment but must consider the unknowns in the overall interpretation of risks and in management decisions (Yoe, 2019, CSAP, 2016). Based on the data gaps and uncertainties summarized in Section 6, the uncertainty in the risk conclusions made to date, and specifically the potential for the assessments to have underestimated risks to human health and the environment, is categorized as moderate to high, and thus, is not supportive of a conclusion of low risk (CSAP, 2016). Rather, based on the uncertainties and the potential for the assessments to have underestimated risk, the risk conclusions are also uncertain, and cannot be further understood until the data gaps are resolved and the uncertainties are decreased.

In the context of biosolids from the CRD, with the data for PFAS and triclosan discussed herein, a review of the data indicates that the concentrations of the COECs are lower than measured in other Canadian biosolids. As noted, the comparison of the low concentrations of COECs with existing toxicity data suggests that the COECs in CRD biosolids represent a negligible to low risk to human health and the environment. Despite this, given the uncertainties discussed in Section 6, the uncertainty in this conclusion is moderate to high and should be reviewed regularly as the science on biosolids evolves.

Oberg and Mason-Renton (2018) examined how uncertainties and gaps in scientific knowledge were addressed and communicated in British Columbia, compared to Sweden, during their jurisdictional review of regulations on the land application of biosolids. The study highlighted how the jurisdictions had approached the uncertainty in the land application of biosolids differently; with BC taking the position that the absence of evidence of risk implies the practice is safe. Sweden, however, prioritized a precautionary approach, operating under the assumption that the absence of evidence or risk is not equivalent to evidence of absence (Oberg and Mason-Renton, 2018). Given the benefits of the land application of biosolids, the Canada-wide approach (CCME, 2012) encouraging the beneficial use of biosolids versus disposal, and as the scientific evidence available to date suggests that the land application of biosolids represents a negligible to low risk, Sweden's approach is likely overly restrictive.

Applying the Precautionary Principle aligns with using an adaptive management framework for the land application of biosolids. As noted, with time, it is anticipated that the uncertainties and data gaps identified in this report and elsewhere will be addressed, and thus regulators must keep pace with the evolving science and regularly weigh the risks and benefits of the practice in an informed and transparent manner. In the interim actions such as source control to limit the introduction of COECs to WWTPs, adopting advanced treatment technologies as they become available, careful site selection through the application of risk-based principles, and ongoing monitoring to minimize risks, are essential for the protection of human health and

the environment. Such strategies are increasingly emphasized in regulatory guidelines around the globe to ensure biosolids use does not compromise human health and the environment. This approach addresses both the potential risks and the benefits of nutrient recycling while minimizing ecological and human health hazards (Schoof & Houkal, 2005; Gianico et al., 2021).

## 9 Limitations

This report has been prepared and the work referred to in this report has been undertaken by Dr. Chris Kennedy for the Capital Regional District (CRD). Dr. Chris Kennedy makes no representation or warranty to any other person with regard to this report and the work referred to in this report and he accepts no duty of care to any other person or any liability or responsibility whatsoever for any losses, expenses, damages, fines, penalties or other harm that may be suffered or incurred by any other person as a result of the use of, reliance on, any decision made or any action taken based on this report or the work referred to in this report.

This report has been prepared based on the CRD Terms of Reference and the literature identified during the review. Dr. Chris Kennedy expresses no warranty with respect to the accuracy of the data reported in the literature.

The evaluation and conclusions reported herein do not preclude the identification of additional literature pertinent to the contaminants discussed in this report. If new literature/studies become available, modifications to the findings, conclusions and recommendations in this report may be necessary.

Where information obtained from reference sources is included in the report, no attempt to verify the reference material was made. Dr. Chris Kennedy expresses no warranty with respect to the toxicity data presented in various references or the validity of the toxicity studies on which it was based. Scientific models employed in the evaluations were selected based on accepted scientific methodologies and practices in common use at the time and are subject to the uncertainties on which they are based.

Nothing in this report is intended to constitute or provide a legal opinion. Dr. Chris Kennedy makes no representation as to the requirements of or compliance with environmental laws, rules, regulations or policies established by federal, provincial or local government bodies. Revisions to the regulatory guidelines and standards referred to in this report may be expected over time, especially considering the evolving nature of the science for many of the contaminants evaluated. As a result, modifications to the findings, conclusions and recommendations in this report may be necessary.

## 10 References

- British Columbia Ministry of Environment and Climate Change Strategy (BC ENV), 2023a. Protocol 1 for Contaminated Sites, Detailed Risk Assessment. Version 4.0. March 20, 2023.
- BC ENV, 2023b. Contaminated Sites Regulation B.C. (BC CSR) Reg. 375/96. OC 1480/96. Last amended March 31, 2023 by BC Reg. 133/2022.
- Behnami, Ali, Khaled Zoroufchi Benis, Mojtaba Pourakbar, Mojtaba Yeganeh, Ali Esrafil, Mitra Gholami, Biosolids, an important route for transporting poly- and perfluoroalkyl substances from wastewater treatment plants into the environment: A systematic review, *Science of The Total Environment*, Volume 925, 2024, 171559, ISSN 0048-9697, <https://doi.org/10.1016/j.scitotenv.2024.171559>.
- Berg C, Crone B, Gullett B, Higuchi M, Krause MJ, Lemieux PM, Martin T, Shields EP, Struble E, Thoma E, Whitehill A. Developing innovative treatment technologies for PFAS-containing wastes. *J Air Waste Manag Assoc*. 2022 Jun;72(6):540-555. doi: 10.1080/10962247.2021.2000903.
- Black GP, Anumol T, Young Thomas M. Analyzing a broader spectrum of endocrine active organic contaminants in sewage sludge with high resolution LC-QTOF-MS suspect screening and QSAR toxicity prediction. *Environ Science: Processes Impacts*. 2019;21(7):1099–114. doi: 10.1039/c9em00144a. [DOI]
- Booton, A., Mayer, B.K. & Zitomer, D.H. Chemical oxidation as an alternative for municipal wastewater secondary treatment: a review. *Rev Environ Sci Biotechnol* **23**, 43–65 (2024). <https://doi.org/10.1007/s11157-024-09684-5>
- Borthakur A, Leonard J, Koutnik VS, Ravi S, Mohanty SK. Inhalation risks of wind-blown dust from biosolid-applied agricultural lands: are they enriched with microplastics and PFAS? *Curr Opin Environ Sci Health*. 2022;25:100309.
- Brown, S., Kennedy, L., Cullington, M., Mihle, A. and Lono-Batura, M. (2019), Relating Pharmaceuticals and Personal Care Products in Biosolids to Home Exposure. *Urban Agriculture & Regional Food Systems*, 4: 1-14 180005. <https://doi.org/10.2134/urbanag2018.12.0005>
- Canadian Council for Ministers of the Environment (CCME), 2012. Canada-Wide Approach for the Management of Wastewater Biosolids. October 11, 2012. PN 1477.
- CCME, 2020. Ecological Risk Assessment Guidance Document. PN: 1585. ISBN: 978-1-77202-044-1 PDF.
- CCME, 2021. Canadian Soil and Groundwater Quality Guidelines for the Protection of Environmental and Human Health. Perfluorooctane Sulfonate (PFOS). Available at <https://ccme.ca/en/res/pfosfactsheeten.pdf>.
- Canadian Food Inspection Agency (CFIA), 2024. Notice to industry: implementation of an interim standard for per- and polyfluoroalkyl substances (PFAS) in commercial biosolids. Available at <https://inspection.canada.ca/en/plant-health/fertilizers/notices-industry/notice-industry-implementation-interim-standard-and-polyfluoroalkyl-substances-pfas-commercial>.
- Cairns, J. Are single species toxicity tests alone adequate for estimating environmental hazard?. *Hydrobiologia* **100**, 47–57 (1983). <https://doi.org/10.1007/BF00027421>
- Carey, Molly. *Fatal Fertilizer: PFAS Contamination of Farmland from Biosolids and Potential Federal Solutions*, 40 Pace Envtl. L. Rev. 282 (2023) DOI: <https://doi.org/10.58948/0738-6206.1870> Available at: <https://digitalcommons.pace.edu/pelr/vol40/iss2/2>
- Chacón, Luz, Liliana Reyes, Luis Rivera-Montero, Kenia Barrantes, Chapter 5 - Transport, fate, and bioavailability of emerging pollutants in soil, sediment, and wastewater treatment plants: potential environmental impacts, Editor(s): Hemen Sarma, Delfina C. Dominguez, Wen-Yee Lee, *Emerging Contaminants in the Environment*, Elsevier, 2022, Pages 111-136, ISBN 9780323851602, <https://doi.org/10.1016/B978-0-323-85160-2.00020-2>.



Clarke, B. O., S. R. Smith (2011): "Review of 'emerging' organic contaminants in biosolids and assessment of international research priorities for the agricultural use of biosolids." *Environ. Int.* 37(1): 226-247

Crossman, Jill, Rachel R. Hurley, Martyn Futter, Luca Nizzetto, Transfer and transport of microplastics from biosolids to agricultural soils and the wider environment, *Science of The Total Environment*, Volume 724, 2020, 138334, ISSN 0048-9697, <https://doi.org/10.1016/j.scitotenv.2020.138334>.

CSAP, 2016. Risk Management Decision Framework for BC Contaminated Sites, Phase 2 – Guiding Principles for Applying Risk-based Standards to Ecological Receptors. Prepared for Society of Contaminated Sites Approved Professionals by Azimuth Consulting Group Partnership.

Duong, H.T., Kadokami, K., Nguyen, D.T. *et al.* Occurrence, potential sources, and risk assessment of pharmaceuticals and personal care products in atmospheric particulate matter in Hanoi, Vietnam. *Environ Sci Pollut Res* **30**, 34814–34826 (2023). <https://doi.org/10.1007/s11356-022-24630-0>

East, A., Egeghy, P.P., Hubal, E.A.C. *et al.* Computational estimates of daily aggregate exposure to PFOA/PFOS from 2011 to 2017 using a basic intake model. *J Expo Sci Environ Epidemiol* **33**, 56–68 (2023). <https://doi.org/10.1038/s41370-021-00374-w>

Eriksen, G. S., C. E. Amundsen, A. Bernhoft, T. Eggen, K. Grave, B. Halling-Sørensen, T. Källqvist, T. Sogn, L. Sverdrup (2009): Risk assessment of contaminants in sewage sludge applied on Norwegian soils. Opinion of the panel on contaminants in the Norwegian Scientific Committee for Food Safety. Norwegian Scientific Committee for Food Safety (VKM). 05/511-22-final.

Evanylo, G., Sukkariyah, B., Anderson Eborall, M., & Zelazny, L. (2006). Bioavailability of Heavy Metals in Biosolids-Amended Soil. *Communications in Soil Science and Plant Analysis*, 37(15–20), 2157–2170. <https://doi.org/10.1080/00103620600817309>

García-Santiago, Xela, Amaya Franco-Uría, Francisco Omil, Juan M. Lema, Risk assessment of persistent pharmaceuticals in biosolids: Dealing with uncertainty, *Journal of Hazardous Materials*, Volume 302, 2016, Pages 72-81, ISSN 0304-3894, <https://doi.org/10.1016/j.jhazmat.2015.09.035>.

Gies EA, LeNoble JL, Noël M, Etemadifar A, Bishay F, Hall ER, Ross PS. Retention of microplastics in a major secondary wastewater treatment plant in Vancouver, Canada. *Mar Pollut Bull.* 2018 Aug;133:553-561. doi: 10.1016/j.marpolbul.2018.06.006.

Goldstein BD. The precautionary principle also applies to public health actions. *Am J Public Health.* 2001 Sep;91(9):1358-61. doi: 10.2105/ajph.91.9.1358. PMID: 11527755; PMCID: PMC1446778.

Government of Canada, 2023. Draft Draft State of Per- and Polyfluoroalkyl Substances (PFAS) Report. Environment and Climate Change Canada and Health Canada. 2023.

Government of Canada, 2024. Updated Draft State of Per- and Polyfluoroalkyl Substances (PFAS) Report. Environment and Climate Change Canada and Health Canada. July 2024.

Grippio, M., J. Hayse, I. Hlohowskyj and K. Picel. 2021. Derivation of PFAS ecological screening values. Final. September 2021. Environmental Science Division, Argonne National Laboratory. Available at [https://www.denix.osd.mil/dodepa/denix-files/sites/85/2022/10/Final-PFAS-ESV-Report\\_Sept-2021\\_508.pdf](https://www.denix.osd.mil/dodepa/denix-files/sites/85/2022/10/Final-PFAS-ESV-Report_Sept-2021_508.pdf).

Health Canada, 2019. Updates to Health Canada Soil Screening values for Perfluoroalkylated Substances (PFAS). May 2019. Prepared by HC Contaminated Sites Division. Ottawa, Ontario.

Health Canada, 2021. Federal Contaminated Site Risk Assessment in Canada. Toxicological Reference Values (TRVs). Version 3.0. ISBN: 978-0-660-36723-1. Pub.: 200301

Health Canada. 2023. Per- and polyfluoroalkyl substances (PFAS) in Canadians. Ottawa, ON. Available: [/content/canadasite/en/health-canada/services/environmental-workplace-health/reports-](https://content.canadasite/en/health-canada/services/environmental-workplace-health/reports-)



publications/environmental-contaminants/human-biomonitoring-resources/per-polyfluoroalkyl-substances-canadians.html.

Health Canada, 2024a. Objective for Canadian Drinking Water Quality, Per- and Polyfluoroalkyl Substances. ISBN: 978-0-660-72710-3, Pub: 240357. Available at <https://www.canada.ca/en/health-canada/services/publications/healthy-living/objective-drinking-water-quality-per-polyfluoroalkyl-substances.html>

Health Canada, 2024b. Federal Contaminated Site Risk Assessment in Canada: Guidance on Human Health and Preliminary Quantitative Risk Assessment (PQRA). Version 4.0. ISBN: 978-0-660-68497-0. Pub.: 230531.

Helmer RW, Reeves DM, Cassidy DP. Per- and Polyfluorinated Alkyl Substances (PFAS) cycling within Michigan: contaminated sites, landfills and wastewater treatment plants. *Water Res.* 2022;210:117983.

Higgins, C. P., J. O. Sharp, J. G. Sepulvado, B. Littrell, G. O'Connor, E. Snyder, D. McAvoy (2010): Trace Organic Chemicals in Biosolids-Amended Soils. State-of-the-Science Review. Water Environment Research Foundation, IWA Publishing. SRSK5T09.

Hill, I.R., Heimbach, F., Leeuwangh, P., & Matthiessen, P. (1994). Freshwater Field Tests for Hazard Assessment of Chemicals (1st ed.). CRC Press. <https://doi.org/10.1201/9780203755488>

Hung W-C, Miao Y, Truong N, Jones A, Mahendra S, Jay J. Tracking antibiotic resistance through the environment near a biosolid spreading ground: resistome changes, distribution, and metal(loid) co-selection. *Sci Total Environ.* 2022;823:153570.

Hydromantis Inc., 2010. Emerging substances of concern in biosolids: Concentrations and effects of treatment processes. Submitted to Canadian Council of Ministers of the Environment, pp. 255.

Jensen, J., S. T. Ingvertsen, J. Magid (2012): Risk evaluation of five groups of persistent organic contaminants in sewage sludge. Danish Ministry of the Environment. Environmental Protection Agency. Environmental Project No. 1406 2012.

Keller, Arturo A., Weiwei Li, Yuki Floyd, James Bae, Kayla Marie Clemens, Eleanor Thomas, Ziwei Han, Adeyemi S. Adeleye, Elimination of microplastics, PFAS, and PPCPs from biosolids via pyrolysis to produce biochar: Feasibility and techno-economic analysis, *Science of The Total Environment*, Volume 947, 2024, 174773, ISSN 048-9697, <https://doi.org/10.1016/j.scitotenv.2024.174773>.

Kennedy/Jenks Consultants. 2017. Biosolids Risk Assessment for Metro Vancouver. Prepared for Metro Vancouver. May 26, 2017.

Kinney, Chad, Brian Vanden Heuvel, Translocation of pharmaceuticals and personal care products after land application of biosolids, *Current Opinion in Environmental Science & Health*, Volume 14, 2020, Pages 23-30, ISSN 2468-5844, <https://doi.org/10.1016/j.coesh.2019.11.004>.

Lakshminarasimman, N., Gewurtz, S.B., Parker, W.J., Smyth, S.A., 2021. Removal and formation of perfluoroalkyl substances in Canadian sludge treatment systems - a mass balance approach. *Sci Total Environ* 754, 142431.

Lavoy M, Crossman J. A novel method for organic matter removal from samples containing microplastics. *Environ Pollut.* 2021 Oct 1;286:117357. doi: 10.1016/j.envpol.2021.117357. Epub 2021 May 14. PMID: 34052645.

Letcher, Robert J, Shaogang Chu, Shirley-Anne Smyth, Side-chain fluorinated polymer surfactants in biosolids from wastewater treatment plants, *Journal of Hazardous Materials*, Volume 388, 2020, 122044, ISSN 0304-3894, <https://doi.org/10.1016/j.jhazmat.2020.122044>.

Marchuk, Serhiy, Stephan Tait, Payel Sinha, Peter Harris, Diogenes L. Antille, Bernadette K. McCabe, Biosolids-derived fertilisers: A review of challenges and opportunities, *Science of The Total Environment*, Volume 875, 2023, 162555, ISSN 0048-9697, <https://doi.org/10.1016/j.scitotenv.2023.162555>.

Mohajerani, Abbas, Bojana Karabatak, Microplastics and pollutants in biosolids have contaminated agricultural soils: An analytical study and a proposal to cease the use of biosolids in farmlands and utilise them in sustainable bricks, *Waste Management*, Volume 107, 2020, Pages 252-265, ISSN 0956-053X, <https://doi.org/10.1016/j.wasman.2020.04.021>.

Oberg (Öberg), Gunilla, Sarah A. Mason-Renton, On the limitation of evidence-based policy: Regulatory narratives and land application of biosolids/sewage sludge in BC, Canada and Sweden, *Environmental Science & Policy*, Volume 84, 2018, Pages 88-96, ISSN 1462-9011, <https://doi.org/10.1016/j.envsci.2018.03.006>.

Pozzebon, E.A., Seifert, L. Emerging environmental health risks associated with the land application of biosolids: a scoping review. *Environ Health* **22**, 57 (2023). <https://doi.org/10.1186/s12940-023-01008-4>

Pramanik BK, Roychand R, Monira S, Bhuiyan M, Jegatheesan V. Fate of road-dust associated microplastics and per- and polyfluorinated substances in stormwater. *Process Saf Environ Prot*. 2020;144:236–41.

Public Health Ontario, 2023. Focus on Per- and Polyfluoroalkyl Substances (PFAS). May 2023. Available at [https://www.publichealthontario.ca/-/media/Documents/P/2023/pfas-per-poly-fluoroalkyl-substances.pdf?rev=270b2dc8401c4653903d3a082579a108&sc\\_lang=en](https://www.publichealthontario.ca/-/media/Documents/P/2023/pfas-per-poly-fluoroalkyl-substances.pdf?rev=270b2dc8401c4653903d3a082579a108&sc_lang=en)

Sivarajah B, Lapen DR, Gewurtz SB, Smyth SA, Provencher JF, Vermaire JC. How many microplastic particles are present in Canadian biosolids? *J Environ Qual*. 2023 Sep-Oct;52(5):1037-1048. doi: 10.1002/jeq2.20497. Epub 2023 Jun 26. PMID: 37296527.

Smith, E.P., Cairns, J. Extrapolation methods for setting ecological standards for water quality: statistical and ecological concerns. *Ecotoxicology* **2**, 203–219 (1993). <https://doi.org/10.1007/BF00116425>

TCEQ, 2021. Chemicals of Interest in Biosolids: Summary of Key Information and Hazard Ranking. February 2021. Available at <https://www.tceq.texas.gov/downloads/toxicology/research-projects/biosolids.pdf>.

Vo, Phong H.N., Gia Ky Le, Lai Nguyen Huy, Lei Zheng, Chawalit Chaiwong, Nam Nhat Nguyen, Hong T.M. Nguyen, Peter J. Ralph, Unnikrishnan Kuzhiumparambil, Soroosh Danaee, Sonja Toft, Craig Madsen, Mikael Kim, Jim Fenstermacher, Ho Truong Nam Hai, Haoran Duan, Ben Tscharke,

Occurrence, spatiotemporal trends, fate, and treatment technologies for microplastics and organic contaminants in biosolids: A review, *Journal of Hazardous Materials*, Volume 466, 2024, 133471, ISSN 0304-3894, <https://doi.org/10.1016/j.jhazmat.2024.133471>.

Wang, Xuebing, Nanyang Yu, Yuli Qian, Wei Shi, Xiaowei Zhang, Jinju Geng, Hongxia Yu, Si Wei, Non-target and suspect screening of per- and polyfluoroalkyl substances in Chinese municipal wastewater treatment plants, *Water Research*, Volume 183, 2020, 115989, ISSN 0043-1354, <https://doi.org/10.1016/j.watres.2020.115989>.

Wang, Yuxin, Kelly D. Good, Microplastics and PFAS air-water interaction and deposition, *Science of The Total Environment*, Volume 954, 2024, 176247, ISSN 0048-9697, <https://doi.org/10.1016/j.scitotenv.2024.176247>.

Warke, Manas, Drew McAvoy, Prioritization of biosolids-borne unregulated organic compounds, *Science of The Total Environment*, Volume 955, 2024, 177207, ISSN 0048-9697, <https://doi.org/10.1016/j.scitotenv.2024.177207>.

WCA, 2019. The Assessment of Organic Contaminants in Material Spread on Land. Final Report to SEPA, from the WCA. January 2019. Available at [https://www.sepa.org.uk/media/413269/organic\\_contaminants\\_materials\\_to\\_land.pdf](https://www.sepa.org.uk/media/413269/organic_contaminants_materials_to_land.pdf).

Xu, B., Liu, F., Cryder, Z., Huang, D., Lu, Z., He, Y., ... Xu, J. (2019). Microplastics in the soil environment:

Occurrence, risks, interactions and fate – A review. *Critical Reviews in Environmental Science and Technology*, 50(21), 2175–2222. <https://doi.org/10.1080/10643389.2019.1694822>

Yoe, C. (2019). *Principles of Risk Analysis: Decision Making Under Uncertainty* (2nd ed.). CRC Press. <https://doi.org/10.1201/9780429021121>

Zhou, Ting, Xuan Li, Huan Liu, Shiman Dong, Zehao Zhang, Zhenyao Wang, Jibin Li, Long D. Nghiem, Stuart J. Khan, Qilin Wang, Occurrence, fate, and remediation for per-and polyfluoroalkyl substances (PFAS) in sewage sludge: A comprehensive review, *Journal of Hazardous Materials*, Volume 466, 2024, 133637, ISSN 0304-3894, <https://doi.org/10.1016/j.jhazmat.2024.133637>.

